



CODE OF PRACTICE



FOR THE CARE AND HANDLING OF DAIRY CATTLE

PUBLISHED 2023

ERRATA ISSUED: JULY 2023 (APPENDIX E)

Canada 

ISBN 978-1-988793-46-7 (book)
ISBN 978-1-988793-47-4 (electronic book text)

Available from:

Dairy Farmers of Canada
45 O'Connor Street, Suite 1410, Ottawa, ON K1P 1A4 CANADA
Website: www.dairyfarmersofcanada.ca
Email: communications@dfc-plc.ca

For information on the Code of Practice development process contact:
National Farm Animal Care Council (NFACC)
Email: nfacc@xplor.net
Website: www.nfacc.ca

Also available in French

© Copyright is jointly held by Dairy Farmers of Canada and the National Farm Animal Care Council (2023).

NFACC is a division of Animal Health Canada (www.animalhealthcanada.ca).

AnimalHealthCanada



This publication may be reproduced for personal or internal use provided that its source is fully acknowledged. However, multiple copy reproduction of this publication in whole or in part for any purpose (including but not limited to resale or redistribution) requires the kind permission of the National Farm Animal Care Council (see www.nfacc.ca for contact information).

Acknowledgment



The project was funded in part by the Government of Canada under the Canadian Agricultural Partnership's AgriAssurance Program.

Disclaimer

Information contained in this publication is subject to periodic review in light of changing practices, government requirements and regulations. No subscriber or reader should act on the basis of any such information without referring to applicable laws and regulations and/or without seeking appropriate professional advice. Although every effort has been made to ensure accuracy, the authors shall not be held responsible for loss or damage caused by errors, omissions, misprints or misinterpretation of the contents hereof. Furthermore, the authors expressly disclaim all and any liability to any person, whether the purchaser of the publication or not, in respect of anything done or omitted, by any such person in reliance on the contents of this publication.



Table of Contents

Preface	4
Introduction	6
Glossary	8
Section 1 Training and Responsibilities	
Section 2 Facilities and Housing	
2.1 Design and Maintenance of Facilities	12
2.2 Housing Systems	12
2.2.1 Calves (Pre-Weaning).....	12
2.2.2 Heifers.....	14
2.2.3 Lactating and Dry Cows	14
2.2.4 Breeding Bulls.....	15
2.3 Facilities for Special Needs.....	15
2.3.1 Calving Areas.....	16
2.3.2 Post-Calving (Fresh Cows)	16
2.3.3 Areas for Sick, Injured, or Lambe Cattle.....	17
2.4 Ventilation, Temperature, and Relative Humidity	17
2.5 Stall Design	18
2.5.1 Electric Trainers in Tie Stalls.....	19
2.6 Space Allowances	20
2.7 Feeding Area	21
2.8 Bedding Management.....	21
2.9 Milking Systems.....	22
2.10 Pasture and Exercise Yards	23
2.11 Emergencies and Safety	23
Section 3 Feed and Water	
3.1 Body Condition Scoring.....	24
3.2 Nutrition and Feeding Management for Cattle	25
3.2.1 Additional Considerations for Heifers.....	26
3.2.2 Additional Considerations for Transition Cows.....	26
3.3 Nutrition and Feeding Management for Calves	27
3.3.1 Additional Considerations for Weaning	28
3.4 Pasture Feeding.....	28
3.5 Water	28
Section 4 Husbandry Practices	
4.1 Handling, Moving, and Restraining Cattle	30
4.1.1 Additional Considerations When Handling or Moving Down Cattle	31
4.2 Surgical and Husbandry Procedures	32
4.2.1 Animal Identification.....	32

4.2.2	Disbudding and Dehorning.....	32
4.2.3	Castration	33
4.2.4	Tail Injuries.....	33
4.2.5	Extra Teat Removal	34
4.3	Udder Hair Removal.....	34
4.4	Breeding.....	34
4.5	Milking.....	35
4.6	Dry-Off Management	35

Section 5 Cattle Health

5.1	Herd Health Management	36
5.1.1	Cattle Cleanliness	37
5.1.2	Pest Control	37
5.2	Genetics.....	37
5.3	Caring for Sick, Injured, or Compromised Cattle.....	38
5.4	Calving Management	38
5.5	Calf Health.....	39
5.5.1	Colostrum.....	40
5.6	Preventing and Treating Mastitis	41
5.7	Promoting Optimal Foot and Leg Health.....	42
5.7.1	Hoof Trimming.....	43

Section 6 Preparations for Transport

6.1	Pre-Transport Decision Making	45
6.1.1	Fitness for Transport (General and Cull Cows).....	45
6.1.2	Additional Considerations for Calves	46
6.1.3	Preparing Cattle for Transport.....	47
6.1.4	Arranging Transport	48
6.2	Loading and Unloading	48

Section 7 Euthanasia

7.1	Decision Making and Criteria for Euthanasia	50
7.2	Methods.....	51
7.3	Confirming Loss of Consciousness and Death.....	52

References..... 54

Appendices:

Appendix A - Sample Cattle Welfare Policy	60
Appendix B - Body Condition Scoring Charts.....	61
Appendix C - Cow Cleanliness Scoring.....	64
Appendix D - Calf Health Scoring Chart and Criteria.....	65
Appendix E - Lameness Scoring System for Dairy Cows.....	67
Appendix F - Transport Decision Tree	69
Appendix G - Sample Euthanasia Decision Tree	70
Appendix H - Anatomical Landmarks for Euthanasia	71
Appendix I - Secondary Steps to Cause Death.....	73

Appendix J - Resources for Further Information.....	75
Appendix K - Participants	79
Appendix L - Summary of Code Requirements	80



Preface

The National Farm Animal Care Council (NFACC) Code development process was followed in the development of this Code of Practice. The *Code of Practice for the Care and Handling of Dairy Cattle* replaces its predecessor developed in 2009 (and published by the National Farm Animal Care Council) effective April 1, 2024 (unless indicated otherwise in Requirements with a later phase-in date).

The Codes of Practice provide critical guidance for the care and handling of farm animals. They serve as our national understanding of animal care requirements and recommended practices. Codes promote sound management and welfare practices for housing, care, transportation, and other animal husbandry practices.

Codes of Practice have been developed for virtually all farmed animal species in Canada. NFACC's website provides access to all currently available Codes (www.nfacc.ca).

The NFACC Code development process aims to:

- link Codes with science
- ensure transparency in the process
- include broad representation from stakeholders
- contribute to improvements in farm animal care
- identify research priorities and encourage work in these priority areas
- write clearly to ensure ease of reading, understanding and implementation
- provide a document that is useful for all stakeholders.

The Codes of Practice are the result of a rigorous Code development process, taking into account the best science available for each species, compiled through an independent peer-reviewed process, along with stakeholder input. The Code development process also takes into account the practical requirements for each species necessary to promote consistent application across Canada and ensure uptake by stakeholders resulting in beneficial animal outcomes. Given their broad use by numerous parties in Canada today, it is important for all to understand how they are intended to be interpreted.

Requirements – These refer to either a regulatory requirement or an industry-imposed expectation outlining acceptable and unacceptable practices and are fundamental obligations relating to the care of animals. Requirements represent a consensus position that these measures, at minimum, are to be implemented by all persons responsible for farm animal care. When included as part of an assessment program, those who fail to implement Requirements may be compelled by industry associations to undertake corrective measures or risk a loss of market options. Requirements also may be enforceable under federal and provincial regulation.

Recommended Practices – Code Recommended Practices may complement a Code's Requirements, promote producer education, and encourage adoption of practices for continual improvement in animal welfare outcomes. Recommended Practices are those that are generally expected to enhance animal welfare outcomes, but failure to implement them does not imply that acceptable standards of animal care are not met.

Broad representation and expertise on each Code Development Committee ensures collaborative Code development. Stakeholder commitment is key to ensure quality animal care standards are established and implemented.



Preface (continued)

This Code represents a consensus amongst diverse stakeholder groups. Consensus results in a decision that everyone agrees advances animal welfare but does not necessarily imply unanimous endorsement of every aspect of the Code. Codes play a central role in Canada's farm animal welfare system as part of a process of continual improvement. As a result, they need to be reviewed and updated regularly. Codes should be reviewed at least every five years following publication and updated at least every ten years.

A key feature of NFACC's Code development process is the Scientific Committee. It is widely accepted that animal welfare codes, guidelines, standards, or legislation should take advantage of the best available research. A Scientific Committee review of priority animal welfare issues for the species being addressed provided valuable information to the Code Development Committee in developing this Code of Practice.

The Scientific Committee report is peer reviewed and publicly available, enhancing the transparency and credibility of the Code.

The Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues developed by the dairy cattle Code of Practice Scientific Committee is available on NFACC's website (www.nfacc.ca).



Introduction

This Code of Practice represents a significant update to the 2009 Code. In updating the Code, the Code Development Committee has endeavored to bring meaningful improvements to animal welfare that can be implemented on the diversity of dairy farms across Canada and that can be defended to producers, customers, consumers, and the public.

This Code is informed by science and promotes the ethical treatment of animals in all situations. This includes ensuring (1):

- good nutrition for normal growth and physiology and to promote satiety and prevent diseases
- housing that is clean and dry, provides thermal comfort, and provides opportunity for exercise and social interaction with other cattle
- disease and injury prevention and prompt detection, followed by appropriate treatments, including nursing care
- positive human-animal interactions that are compassionate and understanding of the animal's emotional needs and that minimize fear and distress
- pain control for painful husbandry procedures.

All dairy cattle deserve to be treated with compassion and respect. Farm owners, managers, and employees have an obligation to provide the same appropriate standard of care to all cattle on their farm irrespective of the animal's economic value.

The committee also recognizes the concept of One Welfare, which holds that animal welfare is interconnected with human wellbeing and the environment. A farmer who is struggling with health or other issues is less able to care for their animals. Likewise, ongoing or sudden animal care challenges on a farm can negatively impact the wellbeing of farmers. The wellbeing of animals and their caregivers is, therefore, interconnected in very important ways.

As a nationally agreed upon standard, this Code of Practice is fundamentally important to improving the welfare of dairy cattle, but, alone, it is not enough. The best outcomes for animals are achieved when a strong national Code of Practice is complemented by on-farm verification to the Code, quality extension and training, and close working relationships between farmers and qualified advisors. The 2009 dairy cattle Code was the basis for the proAction® program, and this updated Code will play a key role in the industry's ongoing efforts to assess animal care on Canadian dairy farms and will guide updates to proAction®.

Requirements in this Code are often outcome- or animal-based, as these are most directly linked to animal welfare and can be applied in a wide range of production systems, giving the producer flexibility to determine how the outcomes can be achieved. Where supported by a body of research evidence, specific outcomes or quantitative criteria have been used.

This Code includes some details specific to the Holstein breed since that is the most common breed on Canadian dairy farms (93%). However, most of the information applies to all breeds of cattle on dairy farms (unless otherwise indicated) and, wherever possible, information unique to other breeds has been included in *Appendix J – Resources for Further Information*.

Scope

In this Code, the word “cattle” refers to dairy cattle of all ages. This Code applies to dairy cattle (including bull calves and bulls) on dairy farms; it also applies to farms dedicated to raising dairy heifers. It does not apply to associated industries (e.g., veal, beef, bull artificial insemination units). Consult the veal cattle and



Introduction (continued)

beef cattle Codes for information on the care of animals in those production systems. For dairy farms that also raise animals for veal, dairy beef, or beef, the dairy cattle Code applies until the animals are moved into separate pens or barns on the farm (or to another farm) dedicated to raising animals for veal, dairy-beef, or beef. The scope of the dairy cattle Code of Practice ends at the farm gate and therefore includes pre-transport Requirements and Recommended Practices.



Glossary

Abusive handling: includes, but is not limited to, kicking, beating, striking, tail twisting, dragging, improper use of a prod, and forcefully pulling cattle by the tail, head, or neck. Refer to additional explanation in *Section 4.1 – Handling, Moving, and Restraining Cattle*.

Ad libitum: allowing animals access to feed or other resources as much as they want when they want (free choice).

All-in/all-out: a production system whereby all animals are moved into and out of facilities and/or between production phases at the same time.

Animal welfare: an animal is in a good state of welfare if (as indicated by scientific evidence) it is healthy, comfortable, well nourished, safe, able to express innate behaviour, and if it is not suffering from unpleasant states such as pain, fear, and distress (2). Animal welfare refers to the state of an animal; the treatment that an animal receives is covered by other terms such as “animal care.”

Calves: male or female bovine animals up to the age at which they are weaned.

Cattle: in this code, “cattle” is used to refer to cattle of all ages.

Cleaning: the process of making equipment or facilities clean by removing wastes such as biofilm, manure, bedding, or other organic debris. (Contrast with “Disinfect.”)

Colostrum: the first milk secreted by the cow after parturition (giving birth), characterized by its high content of proteins and antibodies known as immunoglobulins (Ig).

Competence: demonstrated skill and/or knowledge in a particular topic, practice, or procedure that has been developed through training, experience, or mentorship, or a combination thereof.

Compromised animal: in the context of transport, an animal that has a reduced capacity to withstand transport as indicated by signs of illness, injury, or weakness or due to a specific condition (3) (outlined in *Part XII of the Health of Animals Regulations*). (Contrast with “Unfit animal.”)

Corrective actions: actions to eliminate the cause(s) of nonconformity or other undesirable situations and to prevent recurrence. Generally, corrective actions relate to aspects of animal care or welfare that a producer can control; the action taken needs to be directed at effectively addressing a given issue.

Dehorning: removal of the horns of an animal after the horn buds have attached to the skull. Horn bud attachment occurs at approximately 2 months of age. (Contrast with “Disbudding.”)

Disbudding: removal or destruction of the horn-producing cells of the horn buds before they have attached to the skull (which occurs at approximately 2 months of age). (Contrast with “Dehorning.”)

Disinfection: the application, after thorough cleaning, of procedures or products intended to destroy disease-carrying micro-organisms. (Contrast with “Cleaning.”)

Dystocia: a prolonged calving, which may occur with or without assisted extraction of the calf (4).

Euthanasia: ending of the life of an individual animal in a way that minimizes or eliminates pain and distress (5).

Fit animal: in the context of transport, an animal that is able to withstand the stress of transport without experiencing suffering and is expected to arrive at its final destination in good condition.



Glossary (continued)

Flight zone: in animal handling, flight zone is the space surrounding an animal that, when penetrated, causes the animal to move to re-establish a comfortable distance. Low-stress handling is based on applying and releasing pressure on the edge of the flight zone, ideally never penetrating the zone so aggressively that the animal overreacts and “takes flight.”

Heifers: young female bovine animals from weaning to first calving.

Invasive hoof trim: a trim involving the sensitive laminae of the hoof and/or deeper tissues below the sole, hoof wall, or heel.

Ketosis: a transition period metabolic disorder marked by elevated levels of ketone bodies, indicating that the metabolic processes in the liver are overwhelmed, leading to reduced liver function. Signs include rapid loss of body condition and reduced feed intake (and associated poor rumen fill).

Lameness: any alteration in an animal’s gait that appears to be caused by pain or discomfort. Lameness can manifest as a reluctance or inability to bear weight on a limb, shortened stride, arched back, and/or head bobbing.

Laminitis: inflammation in the digits/hoof that may result in severe pain, abnormal foot growth, and lameness. Laminitis occurs in acute, chronic, or sub-clinical forms, and may be a result of ruminal acidosis.

Local anesthesia or anesthetic: a drug that induces temporary loss of pain sensation in the area to which it is applied.

Low-stress handling: the main principles of low-stress handling are accommodating the animal’s natural behaviours and motivations, reducing noise and other stressors in the environment, and ensuring handlers interact calmly and patiently with cattle.

Neuroma: a mass of regenerating nerve tissue (nerve bundle) that may form when nerve tissue is damaged. Neuromas can result in chronic pain.

Non-ambulatory: unable to rise or stand without assistance or move without being dragged or carried, regardless of size or age (3). An animal must never be dragged (3).

Non-steroidal anti-inflammatory drug (NSAID): a drug that provides pain killing, fever-reducing, and anti-inflammatory effects but is not a steroid or narcotic.

Perinatal mortality: death of a full-term calf at birth or within the first 48 hours.

Personnel: a term used in this Code to refer to any on-farm worker who handles or cares for cattle.

Point of balance: in animal handling, point of balance is the point on an animal’s body (usually the shoulder) where an animal perceives a person to be standing in front of it (causing the animal to back up) or behind it (causing the animal to move forward).

Rumen: the largest of the ruminant stomach chambers and the site of fermentation of fibrous feeds.

Ruminal acidosis: a metabolic condition that occurs when the acidity of the rumen is abnormal (i.e., pH lower than 5.5). The effects can range from disturbance of rumen function (resulting in decreased productivity) to metabolic and health disorders arising from absorption of acids and toxins from the rumen. Acidosis can occur after rapid or over-consumption of highly digestible feeds such as grains.



Glossary (continued)

Rumination: the contractions of the reticulorumen (i.e., the first 2 stomach chambers) to regurgitate previously consumed solid feed, chewing of the feed for a second or more time, followed by swallowing to return the bolus of digesta to the reticulorumen.

Satiety: a feeling of fullness or that hunger has been satisfied.

Sedative: a drug that depresses central nervous system activity, reducing mental activity and body reactions.

Standard operating procedures: written step-by-step instructions describing how a particular task is to be done. Standard operating procedures typically include specific assignment of responsibilities, workflows, desired outcomes, and contingencies.

Systemic analgesia or analgesic: a drug that relieves pain. Systemic analgesia/analgesics provide general (as opposed to local) painkilling effects.

Thermoneutral zone: the range of ambient temperatures at which an animal can maintain a constant body temperature with minimal energy expenditure (6).

Training: a term used in this Code to refer to formal and informal training methods that help ensure personnel carry out procedures correctly and competently.

Udder engorgement: occurs when milk buildup in the udder causes increased intra-mammary pressure, resulting in discomfort and pain. Signs of engorgement can include udder hardness, redness, and heat as well as signs of pain.

Unconsciousness: the point at which an animal can no longer feel pain or perceive and respond to its environment (e.g., light). This state is also referred to as “insensible.”

Unfit animal: in the context of transport, an animal that is unable to withstand transport without experiencing suffering as indicated by signs of illness, injury, or weakness or due to a specific condition (3) (outlined in the *Health of Animals Regulations*). (Contrast with “Compromised animal.”)

Veterinarian-client-patient relationship (VCPR): the basis for interaction among veterinarians, their clients, and their clients’ animals. The VCPR is specifically defined in provincial veterinary acts but, generally, a VCPR has been established when the veterinarian has examined the cattle or visited the farm; the veterinarian has assumed responsibility for making clinical judgments related to the health of the cattle; and the client has indicated a willingness to follow the veterinarian’s instructions.

Veterinary consultation: refers to a one-time consultation or periodic consultations as part of a veterinarian-client-patient relationship. Use of this term is not intended to imply that a consultation with a veterinarian is needed each time the procedure or treatment is carried out.

1

Training and Responsibilities

The people who care for cattle have an important influence on their welfare. Research in several farmed animal species, including dairy cattle, shows that attitudes and beliefs about animals and the importance of routine care influence the way people interact with animals and the diligence with which they carry out their tasks (7). These factors contribute significantly to the variation across farms in productivity and animal welfare outcomes (7).

Farm owners and managers play an important leadership role in ensuring cattle health and welfare are priorities on the farm. Owners and managers also have important responsibilities not only to ensure staff are trained and competent but also to provide ongoing supervision.

REQUIREMENTS

Personnel must be aware of this Code of Practice and must follow the Requirements of this Code of Practice.

Personnel must have the competence to carry out the procedures that they are responsible for.

Managers must supervise personnel and must retrain them if practices begin to fall below standards of care.

RECOMMENDED PRACTICES

- a. develop and implement a cattle welfare policy outlining the farm's commitment to responsible and humane care (refer to *Appendix A – Sample Cattle Welfare Policy*)
- b. identify qualified professionals (on-farm and external) that personnel can connect with regarding cattle care questions or concerns
- c. develop and implement detailed standard operating procedures to support training
- d. review standard operating procedures annually and update them whenever improvements are made to procedures
- e. track training, education, and mentoring activities.

2

Facilities and Housing

Dairy cattle in Canada are housed according to their reproductive state, size, age, and lactation period using a variety of systems. Systems may include loose housing (free stalls, bedded packs, composted packs) or tie stalls, each with or without pasture access. In all life stages, cattle should be housed under conditions conducive to their health, comfort, natural behaviour, and safety. In addition to facility design, management in any given system is always important for ensuring cow comfort.

2.1 Design and Maintenance of Facilities

All housing should be designed and routinely maintained to promote cattle comfort and enable low-stress handling. Experienced handlers who understand how cattle react to noise, light contrast, and shadows will be able to move and restrain cattle more smoothly (8).

REQUIREMENTS

Housing systems, including flooring and other components of housing, must be maintained in good condition to minimize lameness and injury.

Electrified crowd gates must not be used.

RECOMMENDED PRACTICES

- a. consult an experienced agricultural engineer, technical advisors, and other farmers when building a new facility or renovating an existing one
- b. provide soft, high traction flooring in areas where cattle stand for long periods (9)
- c. observe animal walking patterns routinely to assess floors for traction and surface conditions (e.g., level, abrasiveness, obstructions)
- d. minimize the time cows spend on concrete alleyways (9)
- e. flush and/or scrape alleyways and holding areas 2–3 times per day
- f. if cattle balk when moving through facilities, address the reason for balking, which may include increasing the light intensity, ensuring consistent light intensity, or improving flooring
- g. ensure the entrance to a restraint device is well lit and avoid moving cattle from light to dark areas (8)
- h. ensure restraint devices do not exert uncomfortable pressure points on an animal's body.

2.2 Housing Systems

2.2.1 Calves (Pre-Weaning)

Whether housing calves individually, in pairs, or in groups, several management strategies are key to keeping calves healthy and thriving, especially hygiene, clean bedding, air quality, colostrum management, and attentive daily observation to detect the earliest signs of illness.

The stress that calves experience (e.g., from isolation, during weaning) impacts their disease susceptibility, growth, and welfare by decreasing feed intake and by negatively impacting their immune system. Research shows that raising calves in pairs or small groups can reduce stress and improve calf growth, welfare, and learning ability (10). Calves housed with a companion or in a group spend more time feeding, are less fearful, and cope better with novelty (11). Early paired/grouped calves vocalize less during weaning, have healthier eating patterns after weaning, and resume feeding much sooner when moved into larger groups after weaning (12, 13). The behavioural, cognitive, and performance advantages of social housing occur

when calves are paired/grouped early (14, 15, 16) (ideally by 2–3 weeks of age). Calves' strong motivation for social contact (17, 18) is best met by providing full physical contact.

Research results on the impact of social housing on calf health are mixed. While some farms manage large groups successfully, the most consistent finding is a higher occurrence of respiratory disease in larger groups (>8–10 calves) compared to small groups or if air quality is poor (17). No clear trends in calf health (disease incidence, number of treatments, mortality) have been reported in studies comparing individual and group housing methods that have the same milk feeding system and allowance (11). Higher mortality has been found in large groups (>10 calves) compared to individual housing, but no differences have been found between small groups (<7–10 calves) and individual housing (17).

Transitioning to group housing methods necessitates careful planning and implementation to ensure good outcomes. Good management includes evaluating the health status and compatibility of individual calves before they are paired/grouped. Management factors that reduce disease risk in group housing systems include all-in/all-out management, reducing group size (<8–10 calves/pen), and increasing space allowance. Research shows that feeding and weaning strategies can reduce the occurrence of cross sucking (refer to *Section 3.3 – Nutrition and Feeding Management for Calves*).

Hutches are a good housing option when they provide access to outdoor areas and improved air quality (critical for calf health), and permit social interactions by virtue of their design, size, or the way in which they are arranged.

Tethering of calves is not permitted in some Canadian Codes of Practice and has been phased out in other countries. This Code of Practice only allows calves to be tethered to hutches if calves have additional benefits of an outdoor area and therefore fresh air and increased space allowance. However, farmers are strongly encouraged to use hutches that allow calves to have untethered freedom of movement and social interactions to align with the long-term social sustainability of the industry, research on consumer/public viewpoints, and the future direction for dairy cattle housing more generally.

REQUIREMENTS (continued on next page)

For all calf housing systems:

Housing must allow calves to easily stand up, lie down, turn completely around, stand fully upright (without touching the top of the enclosure), adopt sternal and lateral resting postures, groom themselves, and have visual contact with other cattle.

The bedded area for group-housed calves must be large enough to allow all calves to rest comfortably at the same time.

Where tethering of calves is permitted, the tether must include a collar.

For indoor calf housing:

Calves must not be tethered as part of normal indoor housing.

Producers raising calves individually must develop a plan to transition to pair/group housing methods, in consultation with a veterinarian or other qualified advisor.

Effective April 1, 2031, calves that are healthy, thriving, and compatible must be housed in pairs or groups by 4 weeks of age.¹

¹ Movement into pairs/groups may need to be delayed for individual calves that are not healthy and thriving. Once moved into pairs/groups, individual calves may need to be singly housed temporarily if they have a health condition that would improve with separation. Movement into pairs/groups may also need to be delayed to ensure there are sufficient number of calves that are compatible as to their age, size, and drinking speed.

REQUIREMENTS (continued)

Hutches and other outdoor housing:

Calves housed outdoors, including in hutches, must have physical contact with another calf unless they need to be separated for health reasons or they need to be protected from inclement weather.

Calves may be tethered only if housed in hutches that provide access to an area outside the hutch.

RECOMMENDED PRACTICES

- a. select hutches that optimize social interactions (e.g., 2 hutches together with a shared outdoor space, hutches designed for pairs/small groups)
- b. use observations of calf appearance, behaviour, growth, disease, and mortality to evaluate the success of any calf housing system
- c. group calves of similar size and age together to minimize disease risk and competition at feeding
- d. once groups are formed, keep them as stable as possible (introducing a younger calf to an older group, or vice versa, can increase disease risk and competition)
- e. where feasible, manage groups in an all-in/all-out method to minimize disease transmission and permit effective cleaning and disinfection.

2.2.2 Heifers

Having properly designed and managed heifer housing helps ensure well developed replacement animals that will transition well into housing for mature cattle. A growing animal's needs change with age. Flexible systems that accommodate changing requirements in management, housing, feeding, and overall animal care are ideal.

REQUIREMENTS

Housing must allow heifers to easily stand up, lie down, adopt normal resting postures, groom themselves, and have visual and physical contact with other cattle.

RECOMMENDED PRACTICES

- a. provide heifers with daily access to exercise yards or pasture, weather permitting
- b. use housing systems that optimize freedom of movement and social interactions in groups
- c. group heifers of similar size and age together to promote compatibility.

2.2.3 Lactating and Dry Cows

Housing systems should be designed, constructed, operated, and maintained to meet the needs of the cows. There are important advantages and disadvantages to all housing systems—this is reflected in both the research and everyday experience of those who care for cattle. Regular freedom of movement and the ability to socialize are among the benefits of loose housing, but cows can experience competition in these systems. Tie stalls offer cattle a competition-free environment that facilitates observation of individual cattle and, therefore, earlier detection of changes (e.g., body condition). However, freedom of movement is only possible when they're given access to loose housing or an outdoor area.

Research shows no overall benefit for several welfare indicators (e.g., lameness, hock and knee injuries) when comparing tie and free stalls (6, 9). Yet cattle clearly benefit from, and are motivated to have, regular opportunities to move freely (6). Increased movement opportunity through less restrictive indoor housing and/or outdoor access improves dairy cow health, behaviour, performance, and welfare (6). More specifically, giving cattle regular access to open outdoor areas or bedded packs improves hoof health,

reduces the frequency and severity of injuries, and can reduce the occurrence of lameness by 3.5–8% (6). Outdoor access and/or less restrictive indoor housing also enables social grooming and walking/trotting. More research is needed to determine ideal frequencies and durations of freedom of movement, and what constitutes sufficient regular opportunities for freedom of movement will be defined according to research as it becomes available.

The number of tie stall barns being built in Canada and internationally has been steadily declining for many years. Farmers building a new barn are encouraged to continue this trend, one that was initiated within the dairy industry and that aligns with research on consumer/public viewpoints and the long-term social sustainability of the industry. Farmers building a new barn are encouraged to select options that most effectively achieve the Requirement below for daily, untethered freedom of movement and social interactions year-round. Bedded packs and free stalls are among the many examples of systems that effectively meet the needs of the cattle in our care.

REQUIREMENTS

Housing must allow lactating and dry cows to easily stand up, lie down, adopt normal resting postures, groom themselves, and have visual and physical contact with other cattle.

Effective April 1, 2027, cows must not be tethered continuously throughout their entire production cycle (calving to calving) – they must be provided sufficient regular opportunity for freedom of movement to promote good welfare.

Newly built barns must allow daily, untethered freedom of movement and social interactions year-round.

RECOMMENDED PRACTICES

- a. as a guide, provide cows with ~50 hours of outdoor access within any given 4-week period, weather and conditions permitting (19)
- b. build a covered exercise yard, especially if in a high rainfall region.

2.2.4 Breeding Bulls

As with cows and heifers, housing for breeding bulls should be designed, constructed, operated, and maintained to meet the needs of the bulls. Bulls are commonly kept with heifers or cows. However, if kept apart from the herd, bulls should be able to see other cattle to minimize the stress of isolation.

REQUIREMENTS

Housing must allow breeding bulls to easily stand up, lie down, adopt normal resting postures, groom themselves, turn around, and mount safely.

RECOMMENDED PRACTICES

- a. provide bulls a minimum of 18 m² (194 ft²) of pen space
- b. if bulls need to be kept apart from the herd, ensure they have visual contact with other cattle
- c. build secure, sturdy housing
- d. design the bull pen so the bull can be fed, watered, and restrained without anyone entering the pen.

2.3 Facilities for Special Needs

Special needs facilities help ensure personnel can provide for the special needs of calving, transition, and sick, injured, or lame cattle. These areas should maximize comfort, minimize stress and competition, and facilitate nursing care.

REQUIREMENTS

Special needs facilities must include a resting surface with bedding that provides comfort, insulation, dryness, and traction.

2.3.1 Calving Areas

Cows are especially active in the hours before calving, so factors affecting the comfort of the calving area are especially important. A separate calving area, whether for group or individual calving, allows for easier observation and management of individual cows and calves.

Newborn calves are susceptible to disease, making it critically important to maintain clean, dry, and well bedded calving areas.

Ideally, calving areas would be used solely for calving; however, there are times where a pen may need to be used for other special needs cattle. In these circumstances, it is important to maintain pen cleanliness to ensure that disease is not spread between animals.

REQUIREMENTS

Calving areas, whether for group or individual calving, must provide the cow and calf an area that is clean, safe, and separated from the lactating herd, and that provides enough space for the cow to be assisted.

Effective April 1, 2029, cattle on all farms must calve in loose housed maternity pens, yards, or pastures that permit them to turn around.

Newly built barns must allow cows to calve in loose housed maternity pens, yards, or pastures that permit them to turn around.

RECOMMENDED PRACTICES

- a. provide soft, high traction flooring (e.g., soft rubber mats, straw pack)
- b. provide 15 m² (160 ft²) of resting area in individual cow maternity pens
- c. provide 14 m² (150 ft²) per cow of resting area in group calving pens
- d. ensure cows, and especially heifers, are familiar with their calving facilities prior to calving to avoid additional stress around the time of calving
- e. avoid moving or regrouping cows after they have been moved to calving areas
- f. monitor and manage cows in group calving pens for aggressive behaviour
- g. add clean, dry bedding frequently to maternity pens
- h. clean and disinfect maternity pens as often as conditions warrant and with consideration to cow safety and facility design
- i. where feasible, manage group calving pens in an all-in/all-out manner to minimize disease transmission and permit effective cleaning and disinfection (20).

2.3.2 Post-Calving (Fresh Cows)

Fresh cows need special care and a less competitive environment, which may be best achieved in loose housing systems if managed as a separate group. Research on both stall and feed bunk space shows that the effects of overstocking are greatest for transition cows, including first lactation animals introduced into a pen of older cows (6). Refer to *Section 2.6 – Space Allowance* and *Section 2.7 – Feeding Area*.

RECOMMENDED PRACTICES

- a. provide soft, high traction flooring (e.g., soft rubber mats, straw pack).

2.3.3 Areas for Sick, Injured, or Lamé Cattle

Sick, injured, or lamé cattle benefit from being housed in areas that facilitate additional care and treatment and allow them to recuperate without having to compete for feed, water, and lying areas. When ill, cows often separate from herd mates if given the opportunity (21). However, isolation is stressful to cattle, and they should only be segregated when necessary to support their recovery (e.g., prevent injury by herd mates) or minimize transmission of a contagious disease.

Refer also to *Section 5.3 – Caring for Sick, Injured, or Compromised Cattle*.

REQUIREMENTS

Areas must be available to segregate, care for, and treat cattle that are sick, injured, or lamé.

RECOMMENDED PRACTICES

- design or modify facilities to have dedicated areas exclusively for sick, injured, or lamé cattle
- ensure sick pens provide enhanced comfort conducive to recovery (e.g., deep bedding or sand, soft rubber mat, supplemental heat, no drafts)
- ensure convalescing cattle that need to be segregated have visual contact with other cattle
- clean and disinfect sick pens after each use.

2.4 Ventilation, Temperature, and Relative Humidity

Ventilation

Good ventilation, whether natural or mechanical, brings in fresh air and effectively removes dust, airborne pathogens, gases, and excess heat and humidity (6). Dust and ammonia irritate animals' eyes and respiratory tracts and can make cattle more susceptible to respiratory infections. However, definitive thresholds for cattle have not been established, including for ammonia (6).

The risk of pneumonia and other calf diseases can be dramatically reduced through good ventilation (without drafts) and the provision of adequate air space (i.e., at least 6 m³ [212 ft³] per calf up to 6 weeks of age and 10 m³ [353 ft³] per calf up to 12 weeks of age) (22). One of the biggest pneumonia risks for young calves is sharing airspace with older cattle (23).

Temperature and Relative Humidity

Mature dairy cattle are generally able to tolerate low temperatures better than high temperatures (6). When the ambient temperature is above the thermoneutral zone, heat stress occurs because heat load (accumulated both metabolically and from the environment) is higher than the animal's ability to dissipate heat (6). The specific ambient conditions (temperature, humidity) that lead to heat stress vary based on the cow's previous temperature acclimation as well as level of milk production, breed, and other factors (6). High-producing cows are most susceptible to heat stress due to the increased energy demands of milk production (6).

The thermoneutral zone for young calves (up to 3 weeks of age) is 15–25°C (6). Particular attention should be paid to temperatures inside calf hutches, which can far exceed ambient temperatures on hot days (6).

REQUIREMENTS

Facilities, including hutches, must provide cattle with fresh air; prevent the build-up of harmful gases, dust, and moisture; and minimize the risk of heat and cold stress.

RECOMMENDED PRACTICES

- a. consider how many days above or below ideal temperatures for cattle are experienced in a typical year when choosing an appropriate cooling system
- b. monitor cattle routinely for behavioural responses to heat stress, which occur prior to dips in productivity (e.g., increased standing time with shorter lying bouts, panting or increased respiration rate, competition for cooling resources, increased drinking bouts) (6)
- c. ensure good ventilation in milking parlour holding pens
- d. limit reliance on misters and sprayers in high humidity regions (6)
- e. always evaluate air quality, temperature, and speed at cattle level (resting and standing)
- f. if ammonia is ever readily detected by smell, test actual concentrations and take remedial action to ensure it stays below 5–10 ppm
- g. remove manure and soiled bedding from facilities frequently
- h. avoid housing calves in the same air space as older cattle to minimize the occurrence of pneumonia (22)
- i. house calves in well ventilated (but draft free) buildings or in hutches that provide ample fresh air through doors, windows, and top vents
- j. avoid situating calf pens in areas of the barn that tend to be cooler (these conditions are associated with higher disease risk for calves) (24)
- k. avoid exposing cattle to sudden extremes of temperature
- l. design and locate enclosures to take advantage of prevailing summer winds and reduce the amount of solar radiation that enters the barn (barns with an east-west orientation allow less heat from the sun to enter) (6)
- m. ensure appropriate seasonal orientation of hutches (i.e., winter: with the opening facing south to maximize sun exposure; summer: in shaded areas or with the opening facing north to maximize shade)

When facing cold stress:

- n. gradually increase energy intake of calves and heifers in anticipation of cold weather to ensure that growth and weight gain are maintained during periods of cold temperatures
- o. protect cattle from wind and moisture
- p. protect cattle, and especially young calves, from drafts (e.g., build temporary walls/shelters in open-sided barns in the winter)
- q. ensure that the relative humidity inside a housing facility does not exceed 75%
- r. provide calves with clean and dry calf coats in addition to deep bedding (6)

When facing heat stress:

- s. avoid unnecessary procedures or other stressors during the hottest times of the day
- t. provide cattle with access to shaded areas (6)
- u. increase air flow by opening barn doors and vents fully and adding more fans, especially in areas where cattle are misted/sprinkled (e.g., holding areas) (6)
- v. open all vents in hutches or elevate the back of the hutch (by 20 cm [8 in], as a guide) (6)
- w. sprinkle or mist the backs of cattle when they are feeding or otherwise away from resting areas (wetting cattle in resting areas may increase mastitis risk) (6)
- x. modify hutches to have a reflective covering in the summer (6)
- y. choose sidewall heights and overhangs of the roof to reduce sun exposure to stalls, walkways, and bedded packs
- z. use shade cloth when and where needed to minimize the effects of sunlight that enters the barn.

2.5 Stall Design

Stalls that are compatible to the body size of cattle allow natural lying positions that are associated with longer lying times, improved ease of movement, and reduced risk of injurious contact with stall dividers (6). Longer stalls and bed lengths increase lying time and decrease lameness and other injuries (6). Some

research suggests that longer stalls and bed lengths may lead to slightly dirtier stalls, suggesting that enhanced stall cleaning routines may be needed when using longer stalls/beds (6).

Stall dimensions should always be considered relative to the size of the animals, genetic improvements and their effect on size of cattle in the future herd, as well as the behaviour of cattle when using stalls. Stall dimensions (width, brisket boards, neck rail placement) and tie-stall tether length should be set to maximize cow comfort and use of the lying area. Tethers must also be long enough to enable cattle to rest in a head back position.

A guide on evaluating cow comfort, troubleshooting stall configuration problems, and recommended stall configurations is provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Stalls and their components must be compatible with the size of the cattle, minimize lameness and injury, and allow cattle to rest comfortably and rise and lie down with ease.

Tethers or other head restraints must allow cattle to rest in a head back position; stanchions are not permitted.

RECOMMENDED PRACTICES

- a. build and/or modify stalls so that they allow cows to lay comfortably for at least 12 hours per day
- b. build stalls that are sized to the measurements of mature cows to ensure that the design and maintenance will keep cattle clean
- c. build stalls with a lying surface length of at least 1.2 x the height of the cow's rump and a width of 2 x the hip bone width of the cow (and, for tie stalls, add another 15–20 cm [6–8 in] in width, depending on the design of the side dividers) (6)
- d. ensure a tether length of at least 1 meter
- e. modify the stall if cattle are frequently seen standing in the stall but not eating or drinking (25) (see also *Section 2.8 – Bedding Management*)
- f. modify the stall if cattle show abnormal resting or standing postures (e.g., perching, kneeling) (25) (see also *Section 2.8 – Bedding Management*)
- g. modify free stalls (or the ambient areas) that are frequently avoided by cows due to stall design or other disturbances (e.g., ventilation)
- h. observe cattle for abnormal movements as they lie down and rise (e.g., brisk movements, hesitation) (25)
- i. evaluate the time it takes cattle to rise or lie down (cattle should ideally rise in 3–5 seconds and lie down in <5 seconds) (25)
- j. routinely observe the legs of the cows over pressure points for signs of abrasions, swelling, or sores and modify stalls if the rate or severity of injuries increases (6).

2.5.1 Electric Trainers in Tie Stalls

Electric trainers are meant to train cattle in tie stalls to step back to the gutter curb when defecating and urinating so that manure and urine do not fall on the surface of the stall (negatively impacting cattle comfort, cleanliness, and udder health). However, research results are inconsistent and electric trainers may impair hygiene and increase hock and foot injuries, indicating that proper placement is imperative for their effectiveness (6).

Correctly positioned electric trainers can help ensure cattle cleanliness and udder health and do not restrict an animal's ability to show signs of estrus. The correct location above the chine is slightly ahead of the point where the back begins to arch when a cow defecates or urinates.

Electric trainers should be a tool of last resort. On some farms, they can be replaced with modified management strategies (e.g., more frequent cleaning and bedding changes).

REQUIREMENTS

Electric trainers must only be used when needed to train or retrain individual cattle.

Electric trainers must be safe, secure, adjustable, and positioned to enable normal eating, standing, and lying behaviour.

RECOMMENDED PRACTICES

- a. position the trainer to 10 cm (4 in) or higher above the top line of cattle after the training period
- b. for training/retraining, position the trainer about 5 cm (2 in) above the top line of cattle for 24 hours and then raise it again to 10 cm (4 in) above the topline of cattle
- c. have trainers on a timer to ensure they are only on for the minimal time needed
- d. consult an electrician for grounding and installation of trainers.

2.6 Space Allowances

Space allowance is itself an important resource-based measure for animal welfare in loose housing systems, but the ideal density for any given farm should also be considered in the context of relevant outcomes outlined in this Code of Practice (e.g., keeping cattle clean, minimizing lameness).

Free Stalls

Research consistently demonstrates that measures of lying behaviour, most notably lying time, are improved when the availability of lying stalls is increased (6). Free-stall stocking density is often expressed as the number of cows relative to the number of stalls (e.g., 110 cows for 100 stalls, which is equivalent to 1.1 cows per stall). Improvements to lying time and feed bunk access are most pronounced when moving from a high stocking density (e.g., 1.5 cows/stall) to an even stocking density (i.e., 1 cow/stall) (6). However, some benefits are still seen when moving from an even density to a low density (e.g., 0.75 cows/stall) (6). Beyond reducing competition, understocking may allow subordinate cows to avoid lying beside a dominant cow (6).

Bedded Packs

More space is generally needed in bedded packs (compared to free stalls) to achieve similar outcomes related to cleanliness and mastitis management (26). Optimal space allowances range from 7.4–15 m² (80–160 ft²) per animal, or higher, depending on several factors (e.g., climate, bedding, pack management, and breed/size) (26). Based on North American studies, at least 9.3 m² (100 ft²) per cow is needed (26). Higher densities may increase pack compaction and cause excessive moisture (26).

REQUIREMENTS

Stocking density must not exceed 1.2 cows per stall in free stall systems.

Effective April 1, 2027, stocking density must not normally exceed 1.1 cow per stall.²

Effective April 1, 2031, stocking density must not normally exceed 1 cow per stall.²

Resting areas in group pens must provide at least 9.3 m² (100 ft²) per Holstein cow.³

² At any time during or after the 2027 and 2031 transition periods, stocking density can go up to 1.2 cows per stall but only temporarily/intermittently.

³ This minimum required space allowance is based on average weights for large breeds (e.g., Holstein) and will be adjusted for medium and small breeds.

RECOMMENDED PRACTICES

- a. maintain a free stall density of 0.9 cows per stall (6)
- b. reduce stocking density, or modify stall components, if cattle are seen lying outside the stall (refer also to *Section 2.5 – Stall Design* and *Section 2.8 – Bedding Management*)
- c. determine optimal bedded pack space allowance based on overall bedding and pack management and increase space allowance (e.g., to 11–15 m² [120–160 ft²]) if cow cleanliness is suboptimal or cows tend not to rest at the same time.

2.7 Feeding Area

Dairy cattle tend to synchronize their feeding activity and if feeding space is limited (i.e., not all cows can feed at the same time) increased competition may occur, preventing access to feed during peak feeding times (e.g., fresh feed delivery, returning from the milking parlour) (6). Synchronicity driven by milking may be reduced in barns using automatic milking systems, but adequate feeder space is still important (6).

Competition at the feed bunk affects vulnerable cows the most (6). Physical barriers, including head locks and feed stalls, can help reduce competition at the feed bunk and increase feeding time, particularly for subordinate cows (27). Compared to post-and-rail barriers, headlock barriers are associated with fewer displacements, less aggression, and more similar feeding times (6).

In addition to feed space and use of physical barriers, overall feeding management is important for ensuring cattle have good feed access (refer to *Section 3.2 – Nutrition and Feeding Management for Cattle*).

REQUIREMENTS

Provide adequate linear feed bunk space to meet the animals' nutritional needs.

RECOMMENDED PRACTICES

- a. in free stall and bedded-pack barns:
 - provide at least 60 cm (24 in) of linear feed bunk space per lactating cow
 - provide at least 76 cm (30 in) of linear feed bunk space per transition cow (6)
- b. ensure feeding surfaces are smooth and approximately 10–15 cm (4–6 in) higher than the standing area
- c. use physical barriers at the feed bunk
- d. ensure alleys at the feed bunk are at least 4.3 m (14 ft) wide to allow easy movement of cattle
- e. adjust the feed rail if neck injuries occur.

2.8 Bedding Management

Of all possible stall improvements, the provision of large amounts of dry bedding has one of the greatest impacts on cow comfort, lying times, and healing of injuries (6). Bedding is still needed on mattresses and mats—the combination of stall base (including mattresses, mats) and bedding together contribute to the softness and good traction of the stall bed (6). Increasing the depth of bedding not only increases lying times and reduces lameness and other injuries, but it also improves cattle comfort by compensating for hard or abrasive surfaces (6, 9).

Bedding quality, namely dryness, is a key component of cattle comfort (6). Cows and calves consistently show a preference for dry lying surfaces and will spend much more time standing when only wet bedding is available (6). Like mature cattle, calves spend most of their time lying down, and wet or insufficient bedding draws away their body heat. Keeping bedded areas dry also improves cattle cleanliness and reduces lameness, ammonia emissions, and fly infestations (9, 28).

REQUIREMENTS

Cattle must have a resting surface with bedding that provides comfort, insulation, dryness, and traction.

RECOMMENDED PRACTICES

- a. provide generous amounts of clean, dry bedding (the more the better and at least 5 cm [2 in]) to help prevent lameness and promote healing of injuries (6, 9)
- b. incorporate a well-maintained bedding guard into stalls and pens to help keep large quantities of bedding in the animal's area
- c. ensure stalls and pens are routinely bedded and raked out
- d. add new sand and level sand routinely in sand-bedded stalls
- e. routinely observe the legs of cows over pressure points for signs of abrasions, swelling, or sores and increase bedding depth if the rate or severity of injuries increases (6)
- f. increase bedding depth and/or improve bedding management if cattle are frequently seen standing in the stall or pen but not eating/drinking or show abnormal resting or standing postures (e.g., perching, kneeling)
- g. use chopped straw to increase absorption of bedding
- h. replace or top up bedding if your knees get wet in 25 seconds of kneeling in the rest area (an indicator that bedding is too wet)
- i. for stalls and bedded packs: add clean, dry bedding daily
- j. for bedded packs: remove cow patties a few times each day to maintain cow cleanliness
- k. for composted bedded packs: bed as needed depending on climate and other factors and till twice a day to maintain cow cleanliness
- l. in the summer, provide bedding that effectively conducts heat away from cattle (e.g., sand) (6)
- m. in the winter, provide straw bedding (which offers more insulation than other bedding types) and ensure the depth permits cattle, and especially young calves, to nest (5, 28).

2.9 Milking Systems

Minimizing stress on cows in the milking facility is very important. As there is a relationship between the amount of time standing on hard surfaces and lameness, it is desirable to minimize the time cows spend away from feed and water and a comfortable stall.

REQUIREMENTS

Milking equipment must be properly maintained and calibrated.

RECOMMENDED PRACTICES

- a. schedule an annual milking system performance evaluation by a qualified technician
- b. ensure milking areas are uniformly lit and increase light intensity if balking is frequent or light levels do not permit effective inspection of cattle or equipment
- c. monitor cow behaviour during milking and refine equipment if signs of discomfort, fearfulness, or restlessness are observed (20, 29)
- d. ensure the milking facility is designed to minimize the time cows spend away from feed, water, and rest
- e. in robot milker barns: have a low-stress system to prompt any cows that are not leaving the milker in a timely way (thereby preventing access by other cows)
- f. in robot milker barns: have a system for identifying cows in early or peak lactation that are not voluntarily milking on time.

2.10 Pasture and Exercise Yards

Cattle are naturally motivated to access pasture and graze (6). Optimal pasture conditions offer not only a comfortable lying surface but also a cushioned surface with good traction for walking and freedom to move around and graze (6). Outdoor access can also be provided in the form of an exercise yard or a sheltered bedded pack, which offer many of the advantages of pasture (6).

Overall hoof health is generally improved the more cows have access to comfortable footing (e.g., pasture, bedded areas) (9). Giving cows more freedom of movement and opportunities for exercise also improves hoof health by increasing blood flow to the feet and legs (9).

RECOMMENDED PRACTICES

- a. ensure pastures and fences (including electric fences) are safe and properly maintained
- b. ensure cattle are provided with shade and protection from inclement weather when provided outdoor access (e.g., natural or artificial shade in the summer, overhead shelter and/or wind barrier in the winter)
- c. ensure areas chosen for pastures and yards have good drainage
- d. ensure trackways to pastures, outdoor feeding areas, and high-traffic gateways have good drainage and traction.

2.11 Emergencies and Safety

Emergencies may arise and can compromise cattle welfare (e.g., power failure, flooding, disruption of supplies). Pre-planning will assist producers to respond in a timely and effective manner, better providing for the welfare of cattle during these events. Comprehensive resources to support emergency planning have been developed separately from this Code of Practice (refer to examples listed in *Appendix J – Resources for Further Information*).

RECOMMENDED PRACTICES

- a. ensure personnel are familiar with emergency procedures
- b. ensure new or renovated facilities are designed with consideration to emergency procedures (e.g., rapid evacuation of cattle, emergency lighting)
- c. develop a plan for evacuating cattle in the event of an emergency (e.g., transport, alternate facilities for temporary housing)
- d. install an effective alarm system for fire and power failure
- e. consult local fire services on fire prevention and mitigation measures including the correct number of fire extinguishers for all facilities
- f. ensure fire extinguishers are maintained according to manufacturer's instructions and that personnel know where they are located and are trained in their proper use
- g. ensure backup generators are available and functional
- h. employ corrective actions in the event of stray voltage problems
- i. ensure electrical panels are not accessible to cattle.

3

Feed and Water

3.1 Body Condition Scoring

Assessing body condition score (BCS) at first breeding, dry-off, calving, and throughout lactation can help determine whether an optimal nutritional program is in place and help troubleshoot health and fertility problems (30).

Excessively fat and excessively thin cows may experience poor welfare (30). Cows that are too thin (BCS ≤ 2) may be experiencing hunger due to inadequate feed intake, have an underlying health condition, or not have enough body fat reserves to maintain good health or support milk production.

Cows with excessive BCS at calving have poor appetites and lower dry matter intake compared to their thinner counterparts, and this may be a result of the cows' biological drive to return to their natural target BCS of 3 during early lactation (30). High BCS at calving (BCS > 3.25) often results in a rapid loss of body condition following calving (30). Farms that have many over-conditioned dry and lactating cows are, therefore, more likely to have many cows that become too thin after calving as well as a higher occurrence of retained placenta and ketosis. High BCS at calving also significantly increases the risk of dystocia (31).

Over-conditioning is most appropriately addressed through herd-level corrective actions focused on the overall feed program, dry off management, and/or reproductive management. Reducing energy intake of individual cows that are over-conditioned negatively impacts the developing calf and is not a suitable corrective action.

Cows fed even moderate energy diets during the dry period easily over consume relative to their energy requirements. Allowing dry cows to consume excess energy results in many changes typical of excessive BCS, even if cows do not appear to be over conditioned (30). Cows fed a high fibre, controlled energy diet to limit intake to near requirements have a better metabolic profile after calving than cows fed higher energy close-up diets (30). Research has demonstrated that increasing BCS at calving from 3.0 to 3.5 (or from 2.75 to 3.00) results in only a modest increase in milk production (32).

REQUIREMENTS

Corrective action must be taken for cattle at a body condition score of 2 or lower (refer to Appendix B – Body Condition Scoring Charts).

RECOMMENDED PRACTICES

- a. use *Appendix B – Body Condition Scoring Charts* to regularly assess body condition, and keep records of the assessments
- b. aim for the following ideal BCS ranges (30, 32): (a quarter point scale is included in Appendix B)
 - dry off: 2.75–3.25
 - calving: 2.75–3.25
 - growing heifers: 2.75–3.25
- c. take corrective action if more than 15% of the herd is above or below ideal BCS for their production stage (33)
- d. refine feeding and management strategies if $> 3\%$ of cattle are over conditioned at calving (i.e., BCS > 4)
- e. ensure cows in early lactation have a BCS of at least 2.25 (30, 32)
- f. avoid a loss of > 0.5 BCS point in early lactation (32)
- g. keep BCS as consistent as possible throughout the dry period.

3.2 Nutrition and Feeding Management for Cattle

A significant portion of the variability in feed intake and milk yield in herds fed an identical diet is attributable to non-dietary factors such as stocking density and the frequency of feedings and feed push-ups (34, 35). Feeding management, therefore, has a major impact on the health and welfare of cattle. Cattle that are not fed adequately will be hungry and are more likely to have reduced immune function (36).

Strategies to improve feed access include (6, 36):

- reducing stocking density
- increasing the quantity of feed offered
- providing feed ad libitum
- increasing the per animal linear bunk or trough space
- using physical barriers to separate cows as they feed
- increasing the frequency of feedings and feed push-ups.

Increasing the frequency of feedings (to at least twice a day) has also been shown to reduce feed sorting thereby improving the quality of diet cows ingest (37, 38).

Rumination facilitates digestion and stimulates chewing and saliva secretion, which may improve rumen pH and function (39). The amount of time cattle spend ruminating is affected by the characteristics of the diet, health, feed intake, degree of overcrowding, and grouping strategies (39, 40). Rumination is more likely to occur when cows are lying down, making it important to ensure that cattle have comfortable resting areas (40).

Changes in rumination rate are an important early indicator of stress or illness (39). A reduction in rumination often occurs sooner than other indicators (e.g., fever, reduced feed intake, decreased milk yield) (39). Cattle with longer lying and rumination times in the week before calving have higher dry matter intake and milk yield during the first two weeks after calving (39). Shorter rumination time, both prior to and after calving, is associated with an increased risk of metabolic disorders (39). Additional guidance on rumination (and assessing rumen fill) are provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Cattle must have daily access to a palatable ration that meets their nutritional needs, promotes satiety, and maintains body condition, health, and vigour.

RECOMMENDED PRACTICES

- a. establish a feed program for all groups of cattle in consultation with a qualified nutrition advisor
- b. ensure the composition of diets reflects production level, reproductive stage, body size, and environmental temperatures
- c. test nutrient content of feed ingredients used
- d. ensure all rations have been balanced and that all feed components used in the ration are good quality and free of spoilage
- e. offer frequent feedings on a consistent schedule
- f. adopt multiple strategies to minimize competition at feeding (e.g., increase feeding frequency, use physical barriers, increase the quantity of feed offered)
- g. ensure continuous access to feed by frequently pushing up feed in the bunk
- h. ensure provision of fibrous feeds that increase chewing activity or the time it takes to consume the ration (this increases salivary secretions, which helps reduce the risk of acidosis)
- i. consider using automatic sensors for monitoring rumination activity of individual cattle (these are integrated in many commercially available monitoring devices, ear tags, or neck collars) (40).

3.2.1 Additional Considerations for Heifers

The energy requirements of heifers are influenced by their size and growth rate as well as ambient temperatures. Good nutrition, particularly protein, helps ensure adequate frame size, wither and hip heights, and growth rates. Growth rates are an important indicator of the success of heifer feeding strategies, particularly feed access.

Links to breed-specific heifer growth targets are provided in *Appendix J – Resources for Further Information*.

RECOMMENDED PRACTICES

- a. group heifers of similar ages and weights to minimize competition and ensure the quantity fed is appropriate to heifer size
- b. benchmark heifer body weight, wither and hip height, and average daily gains at key stages (e.g., after weaning, before breeding) and refine feeding strategies to achieve ideal targets
- c. aim for average daily gains in post-weaned heifers of 0.6 kg (1.3 lb) per day for small breeds and 0.9 kg (2 lb) per day for large breeds.

3.2.2 Additional Considerations for Transition Cows

Transition cows have increased nutrient demands, and inadequate nutrition during this period can lead to metabolic and infectious diseases (e.g., ketosis, fatty liver, milk fever). These health issues impact negatively on animal welfare and are associated with reduced milk production and reproductive performance as well as early culling (41).

Nutritional approaches to managing milk fever involve monitoring the nutrient balance in the dry cow ration, including the balance of cations (e.g., calcium) and anions (e.g., phosphorus, sulfur) in feed. Forages that are high in potassium (a cation) increase the risk of milk fever (41). Anions promote a more acidic metabolic state (lower blood pH) that is associated with a reduced incidence of milk fever (41).

The degree of reduced feed intake around calving is related to the severity of fatty liver immediately after calving, making management practices that increase feed intake in transition cows critically important (41). However, increasing the energy density of the diets after calving (e.g., feeding additional grain, fat supplements) has not been shown to prevent fatty liver (41). Propylene glycol and rumen-protected choline have been shown to prevent fatty liver and ketosis (41).

RECOMMENDED PRACTICES

- a. track the occurrence of ketosis, fatty liver, and milk fever and consult a qualified nutrition advisor on specific strategies to minimize the occurrence
- b. if the occurrence of milk fever is high: evaluate, with a qualified nutrition advisor, the nutrient balance of feed to determine whether a change is needed (e.g., feeding lower potassium forages, addition of anionic products)
- c. minimize the risk of sub-acute ruminal acidosis from higher concentrate diets by ensuring the diet contains sufficient effective fiber
- d. adjust the ration so that a lower dry matter intake of a high quality, palatable feed is possible, but avoid feeding large amounts of concentrates at one time
- e. feed a diet that minimizes feed sorting
- f. monitor dry matter intake, rumen fill, and body temperature of transition cows
- g. increase concentrate gradually according to appetite (e.g., 0.5–0.7 kg [1.1–1.5 lb] per head per day).

3.3 Nutrition and Feeding Management for Calves

Calves are motivated to consume large volumes of milk (more than eight liters per day for Holsteins) (41). Calves especially benefit from higher milk intakes during the first four weeks of life when their ability to digest solid feed is limited (41). However, calves vary in their intakes and low vitality at birth is among the factors associated with reduced intakes in some calves (42). Feeding a high plane of nutrition using whole milk or high-quality milk replacer is not associated with diarrhea (44).

Non-nutritive sucking (e.g., cross sucking) can occur if calves drink their meal too fast or do not have the ability to express sucking behaviour during and after meals (44). Cross sucking is related to hunger; however, increasing the milk allowance without satisfying the motivation to suck may not help resolve the behaviour (44). Post-meal sucking serves a function for calves in that it promotes digestion and satiety by stimulating the release of digestive hormones (44, 45).

Research shows that the following strategies help reduce the occurrence of cross sucking (44, 45):

- feed through a teat (and consider the use of slow-flow teats)
- increase the number of milk meals while ensuring any given meal satiates the calf
- provide a dry teat after feeding
- manage group feeding systems to reduce competition between calves
- ensure appropriate weaning strategies (refer to *Section 3.3.1 – Additional Considerations for Weaning*).

Providing chopped (not ground) forage, alongside starter, can stimulate greater feed intake and improved rumen development prior to and after weaning, particularly in calves fed higher levels of milk (46).

Refer also to *Section 5.5.1 – Colostrum*. Links to breed-specific calf growth rate targets are provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Calves must receive a diet that promotes satiety and maintains health, growth, and vigour.

Newborn calves must be offered a minimum total daily intake of 15% birth weight (6 L for Holsteins) and from 7–28 days of age must be offered a minimum total daily intake of 20% birth weight (8 L for Holsteins) in milk/milk replacer.⁴

The quantity of milk/milk replacer offered to calves at risk of cold stress must be increased.

RECOMMENDED PRACTICES

- provide milk/milk replacer ad libitum or at least twice daily
- offer milk/milk replacer that is approximately 40°C (104°F); offer acidified milk at approximately 15°C (59°F)
- provide milk/milk replacer via a teat or provide a dry teat after milk feeding to satisfy the calf's motivation to suck
- offer calves at least 9 L of milk/milk replacer daily when ambient environmental temperatures are around 10°C (50°F) and offer at least 10 L daily when temperatures are around 0°C (32°F)
- provide high quality calf starter within the first 7 days of age offering small amounts at each feeding for the first few days (47)
- aim to achieve average gains of 1 kg (2.2 lb) per day and avoid a growth check in the days following birth (41, 44)
- manage group feeding systems to reduce competition between calves (e.g., increase the ratio of teats to calves, use barriers between teats, group calves of similar drinking speed together) (41,44)
- consult with a qualified nutrition advisor about the benefits of providing a limited amount of chopped forage to calves (in addition to regular amounts of palatable starter).

⁴ The amount offered may be reduced for individual calves that are not consistently drinking at this level or for individual calves with health problems.

3.3.1 Additional Considerations for Weaning

Signs of stress during weaning (e.g., cross sucking, vocalizations) can be reduced by weaning gradually and by either adjusting weaning age according to each calf's solid feed intake or by weaning after eight weeks of age (44). Solid feed intake, which increases as calves get older, is an important criterion for when to wean (44). Because cross sucking is partly related to hunger, calves that are better established on solid feed are less likely to cross suck around the time of weaning (45).

Calves that are gradually weaned over at least ten days tend to consume more starter and have better weight gains during weaning and in the immediate post-weaning period compared to abruptly weaned calves and those weaned over four days (44, 48). High preweaning dry matter intake and weight gains have been linked to improved milk production and reproductive outcomes (49).

REQUIREMENTS

Calves must be gradually weaned over a period of at least 5 days, and they must be at least 8 weeks old before weaning is completed (44, 48).

RECOMMENDED PRACTICES

- a. initiate weaning when calves are older than 8 weeks of age or when individual calves are consuming at least 1.4 kg (3 lb) of starter daily for at least 3 consecutive days (44, 47)
- b. wean calves over a period of 10 days or more (44).

3.4 Pasture Feeding

Depending on the region and time of year, cattle may benefit from being fed at pasture. Grazed forage can be an excellent source of feed, particularly for heifers, provided the pasture is properly managed and the diet is supplemented with grain, minerals, and/or other forage sources (based on heifer growth rate and pasture quality) (50).

While high-quality pasture in sufficient quantity can fulfill the nutrient requirements of early dry cows (at least four weeks prior to calving), pasture alone may not provide sufficient energy for some breeds in the close-up period (three weeks prior to calving) (50).

Offering supplemental grain to lactating cows fed at pasture helps ensure greater microbial protein synthesis and increased dry matter intake and is generally needed to maintain body condition (51).

RECOMMENDED PRACTICES

- a. develop a feed program for pastured cattle in consultation with a qualified nutrition advisor (e.g., when supplementation may be needed)
- b. test the quality of forage throughout the growing season so that supplementation accurately corrects for seasonal variations in nutrient content (50).

3.5 Water

All cattle need to have good access to water. Daily water needs depend on many factors, including diet, ambient temperatures, metabolic activity (e.g., pregnant, lactating), and health status (e.g., scours). Water plays an important role in digestive processes—cattle with an insufficient supply of water will limit their solid feed intake. While water intake in young calves fed high volumes of milk/milk replacer tend to be modest, calves with good access to water from birth consume more milk, attain better pre-weaning weights, and may have enhanced ruminal development (and therefore improved nutrient availability for growth) (52).

Water quality affects water consumption (36). Cattle may limit their water intake to the point of dehydration if water contains compounds that diminish palatability (e.g., algae, manure, certain minerals in high concentration) (36). Maintaining clean waterers and periodically testing water quality helps ensure water is safe and palatable.

Additional information on water intakes is provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Watering systems must be clean, and cattle must have access to palatable, clean water in quantities to maintain normal hydration and health, taking into consideration factors such as environmental temperature and diet.

Neither ice nor snow are suitable sources of water.

RECOMMENDED PRACTICES

- a. ensure waterers are at a height that is comfortable for all groups of cattle (e.g., 60–75 cm [24–30 in] for cows in free stalls; 46 cm [18 in] for cows in tie stalls)
- b. provide sufficient drinking space to minimize competition (e.g., 8.9 cm [3.5 in] per lactating cow)
- c. check waterers at least once daily to ensure that they are clean and dispensing water at an ideal flow rate
- d. situate watering points at walkthrough areas (e.g., cross-over alleys)
- e. provide water with a depth of at least 10 cm (4 in) in water troughs
- f. test water quality annually and at high-risk periods (i.e., spring, fall) or whenever problems such as reluctance to drink or reduced feed intake are observed.

4 Husbandry Practices

Husbandry practices are routinely carried out for reasons of management, animal welfare, and human safety (20). As outlined throughout this chapter, options for improving animal welfare in relation to these procedures include using the least invasive method, replacing the procedure with another management strategy, breeding cattle so that they do not require the procedure (e.g., polled genetics, high health traits), and/or providing effective pain control (20).

It is not acceptable to perform painful or stressful procedures or alterations on cattle for cosmetic reasons.

4.1 Handling, Moving, and Restraining Cattle

Human-animal interactions significantly impact cattle welfare and farm productivity (e.g., conception rates, annual milk yields, weight gains) (7). The main principles of low-stress handling are accommodating the animal's natural behaviours and motivations, reducing noise and other stressors in the environment, and ensuring handlers interact calmly and patiently with cattle. Cattle that from a young age are consistently handled using low-stress techniques experience less fear, are less likely to become injured, and will be easier to handle (7, 8).

Abusive handling includes, but is not limited to, kicking, beating, striking, tail twisting,⁵ dragging, improper use of a prod,⁶ and forcefully pulling cattle by the tail, head, or neck.

Proper restraint is important not only for human and animal safety but also facilitates correct application of procedures. Depending on the size of the animal and nature of the procedure, cattle may be safely restrained manually or using a halter, chute, or sedative.

Additional resources on humane handling are provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Personnel must be knowledgeable in cattle behaviour and must only use low-stress techniques in the routine handling of cattle.

Electric prods must not be used for routine handling—they must only be used in extreme situations, such as when an animal's safety is at risk.

Abusive handling is unacceptable.

When restraint is necessary the safest, least stressful restraint must be used.

⁵ Tail twisting is distinct from gently raising the tail to briefly immobilize an animal.

⁶ Improper use of a prod includes repeated use, use on sensitive areas (i.e., udder, belly, face, genitals), use on cattle younger than 3 months of age, or when an animal does not have a clear path to move.

RECOMMENDED PRACTICES

- a. understand the field of vision of cattle and apply the principles of flight zone and point of balance when moving cattle (8)
- b. identify and promptly address barriers to cattle movement (e.g., unusual noises, shadows)
- c. provide sufficient area and a well-lit path for cattle to move in the desired direction (8)
- d. provide high traction flooring (8)
- e. move cattle at a slow walk and in small groups
- f. train cattle to use restraint devices to ensure ease of entry
- g. use a restraint that is ideal for the age of cattle and procedure being done
- h. restrain cattle only for as long as needed to safely carry out the procedure.

4.1.1 Additional Considerations When Handling or Moving Down Cattle

All cattle, but particularly down cattle, need to be handled and moved in a calm, patient, and compassionate manner. Where appropriate, lifting down animals can relieve the pressure on muscles and nerves and is an important strategy to prevent secondary injury associated with prolonged recumbency (53). Cows can only be lifted for a short time (minutes) (53). Detailed guidelines on moving and lifting down cattle are provided in *Appendix J – Resources for Further Information*.

Limited use of an electric prod may be necessary in rare cases to help confirm whether a down animal can rise or if euthanasia needs to be considered. For example, it can help assess nerve damage—lack of response to the prod suggests nerve damage and therefore a poor prognosis. Use of a prod in this context must be done carefully and correctly, if at all. Refer also to *Section 7.1 – Decision Making and Criteria for Euthanasia*.

REQUIREMENTS

Apparatus that are designed to lift, move, and support down cattle must be used according to the manufacturer's specifications.

Hip lifters must only be used to lift an animal for a short duration to help an animal stand on its own—they must never be used to move down cattle.

Down cattle must not be moved by hoisting by chain, dragging, or lifting without adequate body support.

Personnel must not repeatedly encourage a down animal to rise if it has demonstrated it cannot get up or move.

If an electric prod is used, it must be used in consultation with a veterinarian and only applied on the rear flank and upper rear leg (twice at maximum) when absolutely necessary to determine if the animal can rise or if euthanasia needs to be considered.⁷

⁷ In consultation with a veterinarian refers to a one-time consultation or periodic consultations as part of a veterinarian-client-patient relationship. Use of this term is not intended to imply that a consultation with a veterinarian is needed each time the procedure or treatment is carried out.

4.2 Surgical and Husbandry Procedures

Procedures that have the potential to cause pain must be performed in a way that minimizes any pain and stress to the animal (20). For most surgical procedures, including minor procedures, research demonstrates an overall benefit of using both local anesthesia (to prevent acute pain) and a non-steroidal anti-inflammatory drug (NSAID; to reduce post-procedure pain) (4). Pain control is best used pre-emptively (4).

REQUIREMENTS

Surgical procedures must be performed by competent personnel following a method developed in consultation with a veterinarian, including the use of appropriate equipment, pain control, and procedures to minimize the risk of infection and other complications (54, 55).⁷

RECOMMENDED PRACTICES

- select, in consultation with a veterinarian, methods that are the most reliable and least invasive for the size/age of animal
- consider, in consultation with a veterinarian, the inclusion of a sedative to minimize animal stress and ease handling (55).

4.2.1 Animal Identification

Branding is not commonly practiced in the dairy industry and alternative means of identification are accepted by most export markets. Both freeze branding and hot-iron branding cause pain and distress in cattle (41, 56).

REQUIREMENTS

Cattle must not be branded.

RECOMMENDED PRACTICES

- use non-toxic paints for temporary markers
- adjust neck, tail, or leg bands to prevent any discomfort.

4.2.2 Disbudding and Dehorning

Disbudding and dehorning are done for the safety of cattle and personnel. All methods of disbudding and dehorning are painful at any age (41, 56). Animals are easier to handle, heal more quickly, and show lower declines in growth rate when the procedure is performed at younger ages (41, 56). Disbudding (removal of the horn bud before it has attached to the skull) is less invasive than dehorning (horn removal after attachment) (41, 56). Horns typically attach to the skull at approximately two months of age (57). Disbudding may need to be delayed in exceptional circumstances—mainly, for a sick calf or one whose buds are insufficiently developed at two months (to ensure proper application of the procedure).

Proper procedures are important to prevent injury, infection, and re-growth (and subsequent need for a repeat procedure) (41). When using caustic paste, it is important to ensure it does not spread to other animals (in groups or adjacent pens) or to other areas of the treated animal (a particular risk if the calves are outside and it rains).

For all methods, including cautery and caustic paste disbudding, a local anesthetic is necessary to reduce the pain during the procedure and an analgesic is necessary to control longer lasting pain (4, 56).

⁷ In consultation with a veterinarian refers to a one-time consultation or periodic consultations as part of a veterinarian-client-patient relationship. Use of this term is not intended to imply that a consultation with a veterinarian is needed each time the procedure or treatment is carried out.

Breeding cows to polled (genetically hornless) sires results in polled calves and is strongly encouraged as a means of avoiding the need for disbudding/dehorning (20, 41, 56).

REQUIREMENTS

Horn bud removal must be done by 2 months of age (41, 56, 58). Only in exceptional circumstances can individual cattle be dehorned after 2 months of age.

When removing buds or horns, local anesthesia and systemic analgesia must be provided (4, 55, 58).

Banding is not an acceptable method of dehorning (59).

If larger horns must be removed, bleeding must be controlled.

RECOMMENDED PRACTICES

- a. breed cows to polled (genetically hornless) sires to avoid the need for bud/horn removal (20, 58)
- b. aim to disbud calves prior to 4 weeks of age
- c. avoid disbudding day-old calves as it may interfere with colostrum intake
- d. avoid disbudding calves at the same time as other stressors (e.g., sickness, at the same moment as weaning or grouping).

4.2.3 Castration

While not a routine practice in the dairy industry, castration is performed on some farms to prevent unwanted reproduction, reduce aggression towards humans and other cattle, and improve meat quality. All methods of castration cause pain and distress at any age (41, 56). This response can be reduced by using sedatives, anesthetics, and analgesics (41). Animals are easier to handle, heal more quickly, and tend to have a decreased stress response when castrated at a younger age (41).

Factors to consider when selecting a method include the acute pain at the time of the procedure, post-procedure pain (and the extent to which that pain can be managed), rate of wound healing, and stress associated with restraint (56). Methods associated with faster wound healing and fewer complications are preferable, and research suggests that wound healing is fastest with surgical methods (56). The risk of incomplete castration is lowest after surgery, intermediate after the use of a rubber ring, and highest after the use of a Burdizzo/clamp (60).

REQUIREMENTS

If castrating cattle, the procedure must be done as early as possible using local anesthesia and systemic analgesia (4, 55, 60).

RECOMMENDED PRACTICES

- a. avoid castrating calves at the same time as other stressors (e.g., sickness, at the same moment as weaning or grouping).

4.2.4 Tail Injuries

Tails can be broken through interactions with the environment (e.g., scrapers, gates) or by being stepped on by other cattle. Given the frequent activity of the tail, such injuries compromise cattle welfare. Docking may be deemed medically necessary to treat a broken tail in specific individual cases.

Tail docking causes some acute pain and brings the risk of post-operative infection and chronic pain due to neuromas (61, 62).

Routine tail docking is not permitted. It provides no overall advantage in terms of cow cleanliness, udder

health, or milk quality (41). Tail docked cattle may also experience greater discomfort from flies as they are not able to use the tail to control flies (41). Trimming the switch can improve cleanliness and worker comfort.

REQUIREMENTS

Cattle must not be tail docked unless medically necessary for an individual animal, and the procedure must be done using pain control (55).

RECOMMENDED PRACTICES

- a. investigate any tail injury, considering the age of animal affected and the location of the injury, to better understand and promptly address the underlying causes.

4.2.5 Extra Teat Removal

Supernumerary teats (also referred to as extra or sprig teats) may be found as extensions of a primary teat, between the front and rear teats, or behind the rear teats. Supernumerary teats can interfere with milking and create another entry point for bacteria, thereby increasing the risk of infection.

Removing supernumerary teats for purely cosmetic reasons is not ethical or acceptable from an animal welfare standpoint.

REQUIREMENTS

If removing extra teats, they must be removed as early as possible using pain control (55).

RECOMMENDED PRACTICES

- a. provide a local anesthetic to desensitize the teat, in addition to systemic analgesia (4).

4.3 Udder Hair Removal

Unclipped udders accumulate more dirt and make it more difficult to effectively clean the teat and sanitize milking equipment. Flame-clipping (a less time-consuming alternative to electric clipping) involves passing a cool flame under the udder to remove the hair. Even though the flame is cool, proper technique is important to avoid injury.

RECOMMENDED PRACTICES

- a. remove hair from udders on a regular schedule
- b. if using clippers, ensure blades are sharp.

4.4 Breeding

Good breeding management and care of pregnant cows positively impacts on the welfare and future performance of cows and calves as herd replacement animals.

RECOMMENDED PRACTICES

- a. consult a veterinarian on herd reproductive performance and arrange for a veterinarian to perform reproductive examinations
- b. match bull weight and stature to heifer/cow size and physical condition
- c. ensure heifers/cows have achieved adequate stature and are in ideal body condition range prior to breeding (refer to *Section 3.1 – Body Condition Scoring*)
- d. ensure diligent management of cows in heat that are mounting or displaying other injurious behaviours (e.g., temporary separation)
- e. for natural mating, be vigilant about diseases transmitted by natural service and provide secure footing and adequate ceiling height for safe mounting and breeding.

4.5 Milking

Milking should not be a source of stress for cows. Stressors during milking (e.g., novel surroundings, improper handling, loud noise) may cause cows to temporarily “hold back” milk (through a stress induced response that inhibits milk ejection) (29, 63). Better annual milk yields are attained on farms that use low-stress handling techniques during milking (29).

Refer also to *Section 2.9 – Milking Systems*.

RECOMMENDED PRACTICES

- monitor cow behaviour during milking and refine handling techniques or equipment if signs of discomfort, fearfulness, or restlessness are observed (20, 29)
- milk cows at regular, fixed intervals (e.g., same time each day) ensuring lactating cows are not left unmilked or with overly full udders
- achieve a complete milk-out at each milking without overmilking
- avoid painful or stressful procedures (e.g., injections) during milking (29, 63)
- minimize the amount of time cows are standing in holding areas (e.g., maximum 1 hour) to limit time away from feed, water, and rest.

4.6 Dry-Off Management

Dry off needs to be done properly to mitigate the potential welfare concerns associated with abrupt cessation of milking (53). Some research suggests that intermittent milking over a five-day period reduces milk leakage and time anticipating milking (53). Lower-producing cows (<15 kg/d) show less engorgement than higher-producing cows (>25 kg/d) after dry off (64). Reducing milk production of higher-producing cows prior to drying off helps prevent painful udder engorgement and reduce the risk of clinical mastitis (65). Decreasing milking frequency to once a day helps to lower milk production rapidly, without causing pain or discomfort (66).

Refer to *Section 3 – Feed and Water* for general Requirements that also apply during dry off. Refer to *Section 6 – Preparations for Transport* for information about preparing lactating cows for shipping. Additional guidance on dry-off is provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Cows must not be dried off by restricting water.

RECOMMENDED PRACTICES

- dry off cows by gradually (i.e., over at least 5–7 days) reducing milking frequency and shifting to a lower energy and lower protein diet (66)
- offer ad libitum access to a lower energy and lower protein diet throughout the dry-off process (66)
- move end-of-lactation cows out of the lactating herd and into a separate pen or stall, if possible (66).

5

Cattle Health

Good animal health is an integral component of good animal welfare. Health conditions usually cause pain and discomfort, which negatively impact an animal's wellbeing. Good animal welfare, therefore, requires good animal health, and prevention of disease (or treatment/care early in any disease course) is always best for the animals.

5.1 Herd Health Management

Herd health management contributes to animal welfare by providing strategies for disease prevention, rapid diagnosis, and effective treatment. Veterinarians play a key role in implementing changes to improve disease prevention and achieve herd health goals (67). Having a valid veterinarian-client-patient relationship (VCPR) facilitates collaborative decision-making between the producer and veterinarian and is a prerequisite for obtaining some classes of medications.

Having accurate and complete records helps ensure health outcomes are more accurately tracked (68, 69). Research has also shown that producers who keep accurate and detailed health records achieve lower rates of disease (70).

Resources to support herd health management are provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Producers must have a veterinarian-client-patient relationship.

Disease events, treatments, and mortalities (including cause, if known) must be recorded and records must be kept for at least 3 years to track trends in animal health.

Health records must be reviewed with a veterinarian as part of ongoing herd health and disease prevention planning.

RECOMMENDED PRACTICES

- a. consult with a veterinarian when establishing targets for improving herd health outcomes (e.g., lameness, mastitis)
- b. develop protocols in the following priority health management areas:
 - prevention and control of infectious diseases
 - prevention and control of nutritional and metabolic diseases
 - lameness prevention and hoof care
 - mastitis monitoring and control
 - parasite control
 - vaccinations
 - care of down cattle
 - calving management
 - calf health
- c. ensure health protocols include specific treatment protocols, desired outcomes, contingencies, and criteria for when to consult a veterinarian
- d. review health protocols with a veterinarian at least annually or whenever there is a disease outbreak or significant change in health, housing, or management
- e. ensure treatment records include cattle ID, treatment (and reason for), date, and outcome (e.g., recovery, cull, mortality, adverse reaction).

5.1.1 Cattle Cleanliness

Proper bedding and manure management provide animals with a clean, dry, and comfortable environment and better traction when walking or lying down/standing up. The lowest somatic cell counts and incidence of mastitis occur in herds with clean cows and bedding.

Ensuring calves have clean, fresh bedding is among the most important strategies for promoting calf health, particularly prevention of diarrhea and navel infections (28, 71).

REQUIREMENTS

Cattle must be kept clean to minimize disease, maintain udder and hoof health, and promote cattle comfort.

RECOMMENDED PRACTICES

- a. flush and/or scrape alleyways and holding areas 2–3 times a day
- b. clean individual stalls at least twice daily
- c. provide generous amounts of clean, dry bedding
- d. ensure stalls and pens are routinely bedded and raked out
- e. add new sand and level sand routinely in sand-bedded stalls
- f. use cow cleanliness scoring to assess environmental cleanliness and aim for at least 80% of animals in any group with a cleanliness score of 1 or 2 (out of 4; refer to *Appendix C – Cow Cleanliness Scoring*)
- g. brush cows or provide cattle with self-grooming brushes to help keep them clean.

5.1.2 Pest Control

Pests (e.g., insects, rodents) can introduce infectious disease into barns and may cause discomfort to cattle. While pests cannot be fully eliminated, several prevention and management strategies can be implemented to mitigate pest pressure.

RECOMMENDED PRACTICES

- a. ensure housing areas are as free as possible from pests
- b. use humane and environmentally sound methods of pest control (72)
- c. consult a professional pest control specialist annually to review strategies and help assess the effectiveness of pest management strategies
- d. store feed in rodent-proof facilities and containers
- e. promptly remove spilled feeds, particularly wet feeds (e.g., milk, corn silage, haylage)
- f. ensure proper ventilation and screens to minimize fly access
- g. eliminate or reduce the number of places rodents can use for shelter (e.g., clutter, heavy vegetation around buildings)
- h. locate and eliminate insect breeding areas (e.g., wet or soiled bedding, standing water)
- i. clean out (or top up) bedding more frequently in the summer when insect pressure is higher.

5.2 Genetics

There are complex interactions between genetics, husbandry, and environment; however, selection for high productivity has put additional demands on the cow, leading to an increased incidence of disease and higher rates of involuntary culling. Some consulting companies have developed genetic evaluations for several traits in dairy breeds, including functional traits (e.g., calving ability, somatic cell score, conformation traits).

RECOMMENDED PRACTICES

- a. select bulls and cows for traits that contribute to animal health, welfare, and longevity (e.g., calving ability, mastitis resistance, foot and leg conformation, polled).

5.3 Caring for Sick, Injured, or Compromised Cattle

The likelihood of recovery for any sick or injured animal is greatly influenced by the timeliness and quality of nursing care they receive (53). Down cattle need particularly attentive nursing care, including protection from herd mates (or moving them to a safe place in a humane manner), provision of easy access to feed and water, and ensuring that their posture is changed to prevent secondary issues from being recumbent (53). The time and emotional strain associated with caring for a down animal may need to be weighed against the impact it may have on your ability to provide high quality care to other cattle.

Many health conditions cause pain and discomfort or necessitate treatments or procedures that may cause pain. Pain control is most effective when provided early in the course of a disease and prior to a procedure (4). For most surgical procedures, and many painful health conditions, research demonstrates an overall benefit of using both local anesthesia (to prevent acute pain) and an NSAID (to reduce longer lasting pain) (4).

Animal owners are required by law to immediately report the suspicion of a reportable disease (ideally first to the herd veterinarian). Reportable diseases are listed in provincial and federal health acts.

Refer also to *Section 2.3.3 – Areas for Sick, Injured, or Lamé Cattle* and *Section 7 – Euthanasia*.

REQUIREMENTS

Personnel must be able to detect signs of injury, lameness (including abnormalities in gait or mobility), and disease.

Sick, injured, lame, or suffering cattle must receive prompt care appropriate to their condition (including pain control where necessary), and they must be monitored at least twice a day.

Down cattle must have easy access to feed and water and protection from predators, herd mates, and extreme weather (cold, rainfall, direct sunlight), and they must be provided non-slip footing that supports recovery.

Lactating cows that are severely lame or down that require milking (to prevent mammary engorgement) must be milked where they are located.

RECOMMENDED PRACTICES

- establish a system to identify cattle that need to be monitored more frequently due to injury, illness, or other reason
- monitor body temperatures of animals that are sick or being treated
- track changes in weight and/or body condition of sick animals especially calves
- turn down cattle or roll them from side-to-side every few hours if they are not adjusting their own position (or float them for 6–8 hours) to relieve the areas that are bearing weight
- ensure lactating animals that are convalescing are milked at least every 12 hours to relieve udder pressure and reduce the risk of mastitis (53).

5.4 Calving Management

Delivery without complication is the norm in cattle. When a difficult calving (dystocia) occurs, providing timely, high-quality intervention greatly improves outcomes for the cow and calf (31). Dystocia is associated with increased risk of metritis and other reproductive problems, and it is the most important cause of perinatal calf mortality (31). The most important risk factors for dystocia include pre-calving body condition score ≥ 3.5 , first-parity cows, age-at-first calving < 24 months, long gestation length (> 285 days), and heavier calves (31).

The quality of calving management and supervision significantly influences the occurrence of perinatal mortality: mainly, ensuring cows are moved before or after stage 1 of parturition and intervening early in stage 2, if needed (31).

Refer also to *Sections 2.3.1 – Calving Areas* and *5.5 – Calf Health*.

REQUIREMENTS

Steps must be taken to ensure cattle calve in a designated calving area.

Cattle close to calving must be monitored daily, at intervals suitable to individual cattle needs, including for risk factors of dystocia.

RECOMMENDED PRACTICES

- a. develop, in consultation with a veterinarian, calving management protocols (e.g., pre-calving movement, calving location, supervision, intervention, and post-calving management)
- b. ensure heifers have achieved 55–65% of mature body weight at breeding (31)
- c. ensure cows, especially first-parity cows, are at ideal body condition score at calving (2.75–3.25 out of 5) (30, 32)
- d. monitor cattle close to calving every 4 hours and ensure they are checked in the evening and are among the first to be checked in the morning
- e. use cameras to enable frequent, stress-free monitoring of close-up or calving cattle
- f. use automatic sensors to help predict the calving date (e.g., rumination, activity, or body temperature sensors) (31)
- g. ensure cattle are moved before or after stage 1 of parturition (e.g., dilation of the cervix with signs of contractions; moving them during stage 1 interrupts the calving process, which increases the risk of stillbirths) (31)
- h. when cows are first observed in stage 2 of parturition (e.g., feet or amniotic sac are present at the vulva), assess for signs of difficulty and provide timely assistance if needed (e.g., abnormal amniotic fluid; the calf's tongue, head, or feet are swollen or cold) (31)
- i. strive to monitor cows in stage 2 of parturition every 15–30 minutes to confirm delivery is progressing and check for any signs of poor vigour in the calf (31)
- j. consult a veterinarian about difficult calvings and discuss strategies to improve on-farm technique for assisting those cows (73, 74)
- k. consider, in consultation with a veterinarian, providing an NSAID to cows that had a dystocia (4)
- l. clean and disinfect tools/equipment used to assist a calving after each use.

5.5 Calf Health

Calf management in the first days and weeks after birth is integral to setting calves up for success (75). In any calf rearing stage, prompt intervention upon the earliest signs of poor vitality or illness greatly improves calf survival, treatment success, and later productivity (22, 31).

All calves, but especially those born from a dystocia, benefit from attentive care. Calves born from a dystocia are at increased risk of mortality (in the perinatal and later stages) and later health issues (e.g., respiratory disease and diarrhea) (31).

Dipping of calf navels shortly after birth has not been proven to reduce the incidence of navel infection. It can, however, promote drying of the cord and kill any bacteria already present on the surface. If navel dipping is performed, ensure that a clean, 7–10% tincture of iodine is used. Using contaminated solutions or those with additional ingredients such as emollients are detrimental to navel healing.

Fluid therapy (i.e., replacement of lost water and electrolytes) greatly improves outcomes for scouring calves (24). In addition, continuing to offer normal amounts of milk/milk replacer to scouring calves

does not prolong or worsen diarrhea and it prevents weight loss, provides the nutrients necessary for intestinal healing, and supports overall recovery (24, 76). However, feeding milk/milk replacer via tube to sick calves that are not drinking is not recommended as it brings serious health risks.

The threshold for mortality in the Requirement below is informed by research on female calves. While the Requirement focuses on female calves, corrective actions would apply to, and benefit, all calves—male and female.

Refer also to *Appendix D – Calf Health Scoring Chart and Criteria*.

REQUIREMENTS

If mortality in female calves from 2 days of age exceeds 10%, corrective actions must be implemented to improve calving management and calf health in consultation with a veterinarian or other qualified advisor (77).

RECOMMENDED PRACTICES

- a. consult a veterinarian on calf health management as part of routine farm visits (75, 78)
- b. aim to keep all calf mortality below 6% (77)
- c. wherever feasible, designate specific personnel to specialize in calf care
- d. monitor calf body temperature for 2 weeks following birth (normal range is 38.5–39.5°C [101.3–103.1°F])
- e. monitor drinking speed and milk intake in young calves (reductions in these are often an early indicator of illness)
- f. upon the earliest signs of diarrhea, provide fluids (in addition to milk/milk replacer) to replace lost water and electrolytes and promote calf survival (24)

Key strategies for newborn calves:

- g. assess calf vitality as soon as possible after birth (ideally within the first hour) so that timely assistance can be provided if needed (31)
- h. provide additional care or supportive therapy to twins, calves with poor vitality, and those born from a dystocia (e.g., electrolytes, additional colostrum feedings, dry the hair coat and provide additional thermal support)
- i. consider, in consultation with the herd veterinarian, providing an NSAID to calves born from a dystocia (this therapy reduces pain from the delivery and may improve other welfare outcomes) (4).

5.5.1 Colostrum

Colostrum contains antibodies known as immunoglobulins (Ig) that protect calves from infections. Calves' ability to defend against infections is directly related to the amount, quality (at least 50 mg/ml Ig), and timing of colostrum intake (41). Calves' ability to absorb the immunoglobulins from colostrum is substantially reduced six to eight hours after birth (41). Ideally calves are fed using a nipple. However, it is accepted that some calves will receive their first colostrum meals via esophageal tube feeder to facilitate getting the proper volume of colostrum into the calf within the desired time frame. The amount needed depends on the quality of the colostrum. The result of adequate colostrum intake at birth is a high concentration of circulating immunoglobulin in the blood of the calf (i.e., successful passive transfer of immunity) (41).

Calves fed colostrum, transition milk, or a blend of milk and colostrum for the first few days of life have better gut health and development compared to calves that transition to milk after the first colostrum meal (79). Extended colostrum feeding (for the first 2 weeks of life, as an example) has been shown to increase average daily gain, reduce diarrhea (and associated antimicrobial treatments), and improve survival in calves with diarrhea (80, 81).

Additional resources on colostrum management are provided in *Appendix J – Resources for Further Information*.

REQUIREMENTS

Male and female calves must receive at least 4 liters of good quality colostrum within 12 hours of birth, with the first meal occurring as soon as possible, and no later than 6 hours after birth (41).

RECOMMENDED PRACTICES

- a. check the quality of colostrum with a colostrometer or refractometer (41)
- b. measure immunoglobulin status in a sample of male and female calves and strive to ensure all tested calves have a serum total protein concentration of ≥ 5.2 g/dL (82, 83, 84)
- c. consult a veterinarian or qualified advisor if serum protein concentrations are suboptimal
- d. provide supplemental colostrum feeding at birth even when calves are allowed to suckle from the cow (41)
- e. follow strict hygiene practices when collecting, storing, and feeding colostrum (bacterial contamination impedes Ig absorption) (41)
- f. consider feeding male and female calves colostrum, transition milk, or a blend of colostrum and milk for an extended period (e.g., at least 3 days), particularly for unthrifty or low birth weight calves.

5.6 Preventing and Treating Mastitis

Mastitis is a locally painful inflammatory condition of the mammary gland in response to trauma or infection with micro-organisms. Depending on its severity, it can cause systemic illness and associated fever, dehydration, recumbency, and even death. Both local and systemic infection are associated with reduced milk yield and feed intake (85). Mastitis is spread from an infected cow's udder and teat skin to uninfected cows during milking (contagious mastitis) or via micro-organisms (namely bacteria) in the cow's environment that enter the teat ends (environmental mastitis).

Cows with subclinical infections show no overt signs but can still infect other cows. Management of mastitis therefore includes routine somatic cell count testing to identify subclinical cases (threshold of 200,000 somatic cells/ml) in addition to tracking clinical cases. Signs of clinical mastitis include udder abnormalities (e.g., swelling, heat, hardness, redness) and changes in milk (e.g., watery appearance, flakes, clots).

There is a strong body of evidence supporting the use of an NSAID for severe mastitis to reduce inflammation and other indicators of pain (4). Severe cases are those with abnormal milk, with or without udder changes, but with signs of systemic illness such as fever, elevated heart or respiratory rate, dehydration, or decreased rumination (4).

The consistent use of pre- and post-milking teat disinfection procedures combined with proper udder preparation, milking technique, and equipment function are all important factors to preventing udder infection. Keeping the cow and her environment clean and dry is a key factor to reducing the risk of udder infection.

In addition to providing a clean environment, teat sealants can be an important strategy to protect dry cows from environmental mastitis (refer also to *Appendix J – Resources for Further Information*).

REQUIREMENTS

Systemic analgesia must be included in the treatment of cows with severe acute clinical mastitis (4).

RECOMMENDED PRACTICES

- a. consult with a veterinarian to develop a mastitis monitoring and control program using individual cow somatic cell counting and strategic milk culturing
- b. review records of mastitis cases to identify herd-specific risk factors
- c. maintain bulk tank milk somatic cell count below 200,000 cells/ml (86)
- d. strive for a monthly clinical mastitis incidence of $\leq 2/100$ cows (i.e., $< 24\%$ of cows affected/year) (87)

To prevent contagious mastitis infections:

- e. dip each teat of all cows as soon as possible after removal of the milking unit using an approved teat dip
- f. ensure the dip covers the area of the teat skin that had contact with the teat cup liner
- g. ensure infected cows are milked last or separately from uninfected cows (if this is not possible: disinfect the milking unit between uses)
- h. ensure flies are minimized

To prevent environmental mastitis infections:

- i. thoroughly clean and dry teats before milking paying close attention to the teat end
- j. add clean, dry bedding to facilities frequently
- k. maintain a clean, dry environment to promote cow cleanliness
- l. ensure feed is pushed up and available after milking to encourage cows to continue standing, thereby allowing the teat canals to close before cows lie back down (freshly delivered feed promotes longer post-milking standing times) (88, 89).

5.7 Promoting Optimal Foot and Leg Health

Lameness is a painful condition that results in reduced mobility, dry matter intake, and milk production as well as impaired reproduction and early culling (9). Prompt recognition, diagnosis, and early treatment not only minimize the welfare concerns surrounding lameness but also increase the likelihood of recovery and allow cows to produce to their full potential (9).

Education and training are needed to consistently identify injuries and lameness (9). Knowing the actual number of lame or injured cattle on one's farm has been shown to motivate improvements (9). Establishing clear thresholds for improvement is an effective way to focus the farm's efforts towards a goal and track continuous improvement over time. Thresholds can be set by individual farms or industry wide, including through an assurance program.⁸

How cows are housed and managed has a significant impact on the occurrence of lameness and leg injuries (9). Making targeted, manageable improvements that enhance cow comfort and increase resting time can effectively reduce rates of lameness and injuries (e.g., increase bedding quantity, change the stall base, groove crossover alleys) (9).

Research consistently demonstrates that having cows on pasture for even a short period of time significantly reduces the occurrence of lameness and injuries (6, 9). Pasture access can also result in an improvement in gait (in as little as 4 weeks, as an example) particularly if given during the day when cows are more active and motivated to graze (6).

Cattle are considered lame at a gait score of 3 or higher (on the 1–5 scale used in free stalls) and when they have 2 or more signs of lameness using the scoring tool for tie stalls (refer to *Appendix E – Lameness Scoring Systems for Dairy Cows*). Euthanasia should be considered for any lame animal that is not responding to treatment (refer to *Section 7.1 – Decision Making and Criteria for Euthanasia*).

⁸ Industry thresholds are currently established in proAction®. See the zone thresholds in www.dairyfarmers.ca/proaction/resources/overview. Accessed: December 3, 2022.

REQUIREMENTS

Personnel must regularly observe cattle for signs of lameness or leg injuries, to diagnose and treat them quickly.

To minimize lameness and leg injuries, producers must set thresholds for the occurrence of lameness and leg injuries and take corrective actions when the thresholds are exceeded.

RECOMMENDED PRACTICES

- a. participate in gait scoring training and other professional development on lameness prevention
- b. review treatment and hoof trimming records at least annually with a veterinarian or other advisors
- c. aim to keep the occurrence of lameness $\leq 10\%$
- d. aim to keep the occurrence of leg injuries $\leq 10\%$
- e. design and maintain the most comfortable environment possible for cows to maximize resting times (e.g., wide stalls, reduced time standing waiting to be milked, prevent overcrowding) (6, 9)
- f. provide generous amounts of clean, dry bedding (the more the better and at least 5 cm [2 in]) to help prevent lameness and promote healing of injuries (6, 9)
- g. minimize exposure to wet, hard, or slippery floors to ensure cows can move comfortably in indoor environments (9)
- h. remove hazards from floors, yards, and laneways that may cause injury to claws, and eliminate, where possible, steps that cattle climb or descend in their daily activities
- i. aim to examine and treat cattle within 48 hours of when lameness is first detected (9)
- j. give cattle with mild lameness routine daytime access to pasture or deep bedded areas or sand or lime stalls to promote healing (6, 9)
- k. balance the ration to prevent sub-clinical rumen acidosis and associated laminitis
- l. avoid feeding large amounts of concentrate in a single feeding (to reduce the risk of laminitis)

Strategies specific to reducing the occurrence of infectious lameness:

- m. maintain a clean, dry environment to promote cow cleanliness especially cleanliness of the legs (9)
- n. ensure frequent footbathing (at least once a week, as a guide) (9)
- o. ensure footbaths are at least 3 meters long to get ample submersion of each of the cow's feet (9)
- p. replace the footbath solution according to the manufacturer's recommendations and clean the footbath between the solution changes.

5.7.1 Hoof Trimming

Preventative hoof trimming (aiming to maintain correct weight bearing and minimize and prevent lesion development) is a key component of preventing lameness (9). Overgrown hooves are a risk factor for lameness, but the ideal frequency of preventative hoof trimming depends on many factors (9).

Therapeutic trimming can lead to high recovery from lameness; however, recovery is dependent on the severity of the lameness with better outcomes reported in lower severity cases (9). Early identification of lameness is, therefore, critically important for improving outcomes (9). Strategies to relieve pain and pressure on the injured area and promote healing include a trim that effectively relieves pressure, a hoof block, or an analgesic.

Improper disinfection of hoof trimming equipment is among the many risk factors for digital dermatitis, and specific disinfectants are needed to kill the associated bacteria (which can survive for several hours on trimming equipment) (9).

REQUIREMENTS

Feet and claws must be inspected and hooves trimmed as required to promote a normal gait and minimize lameness.

Infectious hoof lesions must be treated to control the infection.

Therapeutic hoof trimming must include strategies to relieve pain and pressure on the injured area and promote healing.

Pain control must be included in the treatment of cattle that receive an invasive hoof trim.

RECOMMENDED PRACTICES

- a. examine hooves regularly and trim at least twice a year (90)
- b. avoid preventative hoof trimming in the first month post calving
- c. ensure personnel responsible for hoof trimming are certified and/or affiliated with a professional association
- d. ensure strict biosecurity protocols for hoof trimming particularly cleaning and disinfection of tools between farms and after treating cattle with contagious hoof conditions
- e. ensure accurate and complete hoof trimming records are kept (e.g., cattle ID, date, lesion, treatment)
- f. monitor outcomes following hoof trimming and refine techniques or identify other hoof trimmers if outcomes are suboptimal.

6

Preparations for Transport

Humane transport of dairy cattle is important for their welfare and is expected by the dairy industry and consumers alike. Marketing healthy, sound calves and cows is an important achievement for the dairy producer—all producers should strive to consistently achieve this. Cattle that are in good health and condition prior to transport have better outcomes during and after transport.

The federal requirements for animal transport are covered under the *Health of Animals Regulations, Part XII*.⁹ They are enforced by the Canadian Food Inspection Agency (CFIA) with the assistance of other federal, provincial, and territorial authorities. Some provinces have additional animal transport regulations. Anyone with responsibilities during any part of the transport process (i.e., loading, confinement, transport, or unloading) must be familiar with, and follow, all applicable requirements. Refer to Appendix J for resources to support compliance with transport regulations.

6.1 Pre-Transport Decision Making

6.1.1 Fitness for Transport (General and Cull Cows)

An animal's fitness for transport must be assessed by considering the following risk factors that could impact on their capacity to handle loading, confinement, transport, and unloading (3):

- the current condition of the animal and any pre-existing weakness, disease, or injury
- the space requirements for the animal
- compatibility with other animals
- animal handling and restraint methods
- the expected time without feed, water, and rest
- the expected duration of transport (including intermediate stops, auction markets, and how far they may be transported after the auction)
- the foreseeable delays during transport and at the destination
- the foreseeable weather and other conditions of the journey (e.g., sharp inclines, swaying of the conveyance)
- the type and condition of the conveyance and loading equipment.

If in doubt, assume the longest trip when assessing an animal's fitness. Producers have a primary responsibility for determining if an animal is fit for transport. While the carrier or the driver should not be relied upon to determine whether an animal is compromised or unfit, they have the right and responsibility to ensure special provisions for a compromised animal or to refuse to load an animal that they deem unfit.

Many cull dairy cows are compromised or unfit, and research demonstrates that gait score, body condition, and udder health tend to deteriorate the longer an animal is transported or in the auction-slaughter system (53). Screening for lameness, adequate body condition, and risk of udder engorgement prior to loading is particularly important (53). Transport is not an acceptable solution for a lame or very thin animal. These cattle need on-farm treatment, additional feeding, and/or convalescence before they can be removed from the herd (refer to *Section 3 – Feed and Water* and *Section 5 – Cattle Health*). Euthanasia may be the most humane option for some animals with a condition impacting their fitness (refer to *Section 7 – Euthanasia*).

⁹ The *Health of Animals Regulations* are accessible through the following link: www.laws-lois.justice.gc.ca/eng/regulations/c.r.c.,_c._296/page-10.html#h-548075. Accessed: November 16, 2022.

REQUIREMENTS

Every animal's fitness for transport must be assessed before they are loaded taking into consideration their condition and risk factors that may impact the animal's capacity to withstand loading, transport, and unloading (3).

Unfit animals must not be transported except for veterinary care on the advice of a veterinarian and with special provisions (refer to the regulatory guidance in Appendix F – Transport Decision Tree) (3).¹⁰

Compromised animals must only be transported with special provisions and directly to the nearest suitable place where they can receive care or be promptly slaughtered (not through an auction or assembly yard; refer to the regulatory guidance in Appendix F – Transport Decision Tree) (3).¹⁰

RECOMMENDED PRACTICES

- a. increase the frequency of health monitoring as the shipping date approaches to ensure early detection of a condition that may warrant early shipping or treatment before cattle become compromised
- b. if in doubt about whether an animal can withstand the same transport challenges as a healthy, fit animal, assume the animal is compromised (and transport with special provisions) or consult a veterinarian (91)
- c. when consulting a veterinarian about an animal's fitness for transport, use pictures or videos of the animal to facilitate the consultation
- d. identify local options for slaughter or use mobile slaughter (if available) for cull dairy cows to eliminate or reduce time in transit
- e. consider on-farm emergency slaughter where permitted and appropriate for the animal's condition (e.g., injury preventing transport but no systemic illness, as assessed by a veterinarian).

6.1.2 Additional Considerations for Calves

The care calves receive on the dairy farm greatly impacts their fitness for transport and later health and welfare outcomes. Calves that at the time of shipping are bright and alert; have consumed sufficient, high-quality colostrum; are heavier; and have a healed navel have lower risk of mortality and illness and better growth rates in the weeks following transport (92, 93, 94, 95).

Calves are unfit if they have an infected navel or signs of a fever, dehydration, or exhaustion (all other conditions are listed in *Appendix F – Transport Decision Tree*). Signs of fever include (91):

- body temperature >39.5°C (>103.1°F)
- hot to the touch, particularly in hairless areas
- red tinge to the skin
- panting
- lethargic, weak, or dull/depressed.

¹⁰ The *Health of Animals Regulations* require that compromised or unfit animals are individually loaded and unloaded without having to negotiate ramps inside the conveyance, are isolated during transport (compromised animals may be segregated with one familiar animal), and other measures are taken as necessary to prevent suffering or injury (e.g., mitigate pain, protect from cold, prevent dehydration).

REQUIREMENTS

Calves must only be shipped if they are free from signs of fever or disease and have a healed, uninfected navel (refer to other regulatory guidance in Appendix F – Transport Decision Tree).¹¹

Calves that are 8 days of age or less must only be transported with special provisions and directly to their final destination (not through an auction or assembly yard) (3).¹²

RECOMMENDED PRACTICES

- check the body temperature of calves to confirm they have a normal body temperature (38.5–39.5°C [101.3–103.1°F])
- palpate the navel (consider wearing gloves for hygiene) to ensure it is dry and free of signs of infection (e.g., pink/red, swelling, hot, pain indicators, discharge)
- ship calves that are at least 45 kg (100 lb) and avoid shipping calves that are ≤27 kg (≤60 lb)
- avoid shipping calves that are less than 8 days old
- aim to ship all calves (i.e., including those older than 8 days) directly to the nearest calf grower to reduce handling events and time away from a comfortable pen
- when marketing calves directly to growers: establish a calf purchase agreement detailing colostrum and milk feeding practices, herd and calf health status, weight range, and other conditions for calf guarantee
- wherever feasible for the facilities: market calves directly to growers as pre-conditioned (i.e., weaned, vaccinated, and 80–136 kg [176–300 lb]).

6.1.3 Preparing Cattle for Transport

Preparing cattle for transport starts long before the trip begins. For mature cattle, management factors such as opportunities for exercise, lameness prevention, and nutrition have a collective impact on fitness for transport. Drying cows off, resting them in a comfortable pen, and feeding them to put weight on has the potential to both improve cow welfare and to add value to cull dairy cows prior to transport (53). While cows do not have to be dried off before transport, they must reach their final destination (or a suitable place where they can be milked) before their udders become engorged (91). Refer also to *Section 4.6 – Dry-Off Management*.

Strategies that help prepare calves for transport include good calving management; timely intake of sufficient, high-quality colostrum; provision of ample milk/milk replacer; and keeping calf facilities clean and well bedded (78, 93).

Maximum intervals of time cattle may be without feed, water, and rest are outlined in the *Health of Animals Regulations*. The interval without feed and water begins when animals are last fed and watered on the farm. For fit cattle, the interval without feed, water, and rest must not exceed 36 hours; for compromised cattle, the interval must not exceed 12 hours (3). For calves, the time without feed, water, and rest must not exceed 12 hours (3).

REQUIREMENTS

When shipping a lactating cow, steps must be taken to prevent the risk of udder engorgement.

¹¹ From the *Health of Animals Regulations*: calves with signs of fever or that have an unhealed or infected navel are unfit and must only be transported for veterinary care on the advice of a veterinarian with special provisions.

¹² The *Health of Animals Regulations* require that calves that are 8 days of age or less are individually loaded and unloaded without having to negotiate ramps inside the conveyance; have sufficient space to allow the animal to lie down without lying on top of another animal; are segregated from older animals (other than their dam); and that other measures are taken as necessary to prevent suffering, injury, or death.

RECOMMENDED PRACTICES

- a. ensure good communication between personnel who care for cattle to be shipped and those that will make the decision to ship
- b. ensure good communication with the transporter especially at the time of loading
- c. provide unweaned calves with at least half of that day's ration of milk prior to loading
- d. milk out lactating cows immediately before shipping if the cow is still producing milk on the day of transport
- e. gradually dry off all lactating cows prior to shipping to slaughter (starting at least 5–7 days prior to the shipping date wherever possible)
- f. if shipping a lactating cow to slaughter, aim to ensure they are producing <20 litres/day.

6.1.4 Arranging Transport

Producers have a responsibility to ensure that the transporter they hire is trained and qualified. Producers arranging for transport should also be aware of additional services that may be required under the *Health of Animals Regulations* during transit (e.g., feed, water, rest, milking).

REQUIREMENTS

Personnel involved in loading, unloading, or transporting cattle must have the necessary knowledge and skills to conduct these activities in compliance with the Health of Animals Regulations (3).

RECOMMENDED PRACTICES

- a. schedule transport such that delays are avoided due to severe weather, road construction, or ferry cancellations
- b. ensure all required documentation is completed prior to loading to avoid delays at inspection stations or other checkpoints, slaughter plants, or for cattle leaving the country
- c. ensure the following information is discussed and agreed upon with the transporter:
 - number of cattle to be shipped
 - type of cattle (e.g., yearling heifers, mature cows, bulls, calves)
 - type and quantity of bedding
 - time and point of loading
 - timing of the last provision of feed and water
 - destination
 - special requirements, if any, of the animals being transported
 - special protection from hot or cold weather for all cattle, especially calves
- d. hire transporters that have extensive experience in transporting dairy cattle.

6.2 Loading and Unloading

Using low-stress handling techniques and well-designed handling systems facilitate animal movement and reduce stress and injury. Round crowd pens and curved single file chutes work better than straight ones. Curved cattle chutes are better for handling cattle because they take advantage of the natural tendency of cattle to go back to where they came from (96). Refer to *Section 4.1 – Handling, Moving, and Restraining Cattle*.

Cattle may be considered incompatible for transport based on factors such as their size, weight, temperament, or age. Steps must be taken to ensure incompatible cattle are not transported together. Strategies include avoiding groupings of cattle that will lead to fighting and injury and segregating individual or groups of calves from older animals (other than their dam).

REQUIREMENTS

Ramps, gangways, chutes, and steps must be designed, constructed, maintained, and used in a way that prevents the animal from tripping, slipping, or falling (3).

The slope of ramps used to load or unload animals onto/from the conveyance must not exceed 25 degrees (3).

Cattle that are incompatible must be segregated (3).

RECOMMENDED PRACTICES

- a. use curved chutes with solid sides
- b. ensure loading facilities are uniform in appearance, designed to minimize noise, and uniformly lit (address any sharp contrasts and shadows)
- c. during hot weather, avoid loading during the hottest part of the day
- d. provide overhead cover in loading/unloading areas to protect cattle from wet conditions
- e. minimize handling and non-essential procedures on the day of loading or arrival to avoid exposing animals to stressors in addition to transport.

7

Euthanasia

Euthanasia is necessary when a calf, heifer, cow, or bull is not responding to treatment, medical care to alleviate suffering or pain is not feasible, or there is no reasonable prospect for recovery. Euthanasia may also be necessary for disease control or in a situation where there is a potential for animal suffering arising from a lack of marketing options.

Producers, personnel, and veterinarians who perform euthanasia should be aware that they may be at risk of emotional stress and should take measures to mitigate this risk (97). The impacts on mental health may be more significant for those who have been directly involved in the care of the animals or when several animals need to be euthanized. Where feasible, it may be beneficial to arrange for euthanasia to be performed by someone other than the person who cared for the animal(s).

7.1 Decision Making and Criteria for Euthanasia

Personnel who are responsible for euthanizing cattle play an important role in their welfare. The successful application of any method depends on many factors, particularly the experience, training, and compassion of the individual carrying out the procedure (97).

Timeliness of euthanasia is also critically important to avoid prolonged suffering (5). When someone has worked tirelessly to improve an animal's condition, it can be hard to accept that the animal is not improving (5). However, it is important to recognize when euthanasia is the most humane and responsible course of action in the best interest of an animal.

Euthanasia may be the most humane option for cattle with severe or chronic lameness or when the animal is not responding to treatment.

While down cattle need to be individually evaluated, generally, the longer an animal remains recumbent the less likely they are to recover (53). Cows that are recumbent for more than 24 hours have a poor prognosis, and recovery may be influenced more by secondary factors from being down than by the original primary condition (53).

The likelihood of recovery for any animal, including down cattle, can be influenced by the quality of nursing care (53). Refer to *Section 5.3 – Caring for Sick, Injured, or Compromised Cattle*. A sample euthanasia decision tree is provided in Appendix G.

REQUIREMENTS

Cattle must be promptly euthanized if they have a condition that compromises their welfare AND at least 1 of the following applies:

- ***they do not have a reasonable prospect of improvement,***
- ***they are not responding to treatment(s) within an appropriate timeframe for the condition, or***
- ***treatment is not a humane option.***

Personnel who perform euthanasia must be trained and competent in all aspects of the acceptable method(s) used on the farm.

RECOMMENDED PRACTICES

- a. work with a veterinarian to develop and implement a euthanasia plan to facilitate timely and humane euthanasia
- b. participate in ongoing euthanasia training (e.g., workshops, online videos, peer or veterinary demonstration on farm)
- c. strongly consider euthanasia for any animal down for a prolonged period (e.g., >24 h) (53)
- d. strongly consider euthanasia for any down animal that is:
 - not eating
 - not alert
 - not showing daily improvement
 - not willing/able to lay in a sternal position, or that
 - has pressure sores or any other complication.

7.2

Methods

Euthanasia must be quick, cause minimal stress and pain, and result in rapid loss of consciousness followed by death without the animal regaining consciousness (97). Methods that meet these criteria are outlined in *Table 7.1 – Acceptable Euthanasia Methods for Cattle*. Additional guidance on these acceptable methods is provided in *Appendix H – Anatomical Landmarks for Euthanasia* and *Appendix I – Secondary Steps to Cause Death*. Methods not listed in Table 7.1 are not acceptable.

Proper storage and maintenance of equipment is essential to ensure the equipment functions effectively, particularly with captive bolt guns (5). Proper restraint of the animal is often needed to carry out the method correctly. Humane handling is also an important component of euthanasia (5). It may be preferable to perform euthanasia out of sight and smell of other cattle; however, cattle to be euthanized must only be moved or handled humanely (refer to *Section 4.1 – Handling, Moving, and Restraining Cattle* and *Section 4.1.1 – Additional Considerations When Handling or Moving Down Cattle*) and only if necessary.

REQUIREMENTS

An acceptable method for euthanizing cattle must be used (acceptable methods are listed in Table 7.1).

The method of euthanasia must be quick, cause minimal stress and pain, and result in rapid loss of consciousness followed by death without the animal regaining consciousness.

Manual blunt force trauma is not an acceptable means of euthanasia, including for calves at any age (5, 100, 101).

Every farm must have the ability to euthanize animals or have timely access to euthanasia services.

Equipment necessary for euthanasia (including a secondary step, if applicable) must be used, stored, and maintained according to the manufacturer's instructions to ensure proper function.

Prior to being euthanized, cattle must only be handled or moved if necessary.

When restraint is necessary, euthanasia must be performed without delay following restraint.

RECOMMENDED PRACTICES

- a. consult with a veterinarian when selecting acceptable method(s) of euthanasia
- b. consider, in consultation with a veterinarian, sedation as a means of humane restraint for euthanasia.

Table 7.1 Acceptable Euthanasia Methods for Cattle (adapted from 5, 98, 99, 100)

Method	Suitable for	Equipment and Procedure
Gunshot with a rifle or shotgun	Calves less than 181 kg (400 lb)	<p>Applied to the correct frontal landmark (refer to <i>Appendix H – Anatomical Landmarks for Euthanasia</i>)</p> <p>Examples of appropriate firearms:</p> <ul style="list-style-type: none"> • Rifles: a .22 long rifle fired from a short distance is acceptable (i.e., 0.60–0.90 m [2–3 ft]) • A .22 magnum or larger calibre is recommended • Shotguns: a .410 shotgun with a 7.6 cm (3 in) magnum load with bird shot or slug fired a short distance or a 12 or 20 gauge shotgun with slugs or bird shot no. 2, 4, or 6.
	Cattle more than 181 kg (400 lb)	<ul style="list-style-type: none"> • Requires a minimum of 1356 J (1000 ft-lb) of muzzle energy. <p>Examples of appropriate firearms:</p> <ul style="list-style-type: none"> • Rifles: a .22 magnum or larger calibre centre fire rifle (.223, .270, 303, 30-30) is required • Shotguns: a 12 or 20 gauge shotgun with slugs or bird shot no. 2, 4, or 6. <p>Note: A standard .22 calibre long rifle only produces 119–138 joules (116–135 ft-lb) of muzzle energy and is not sufficient for cattle at this weight.</p>
Drugs approved for euthanasia	All weights and ages	<ul style="list-style-type: none"> • Must be administered by a veterinarian. Safe disposal of the carcass is required when barbiturates are used.
Penetrating captive bolt and secondary step to cause death	All weights and ages	<ul style="list-style-type: none"> • Bolt must be applied to the correct frontal landmark (refer to <i>Appendix H – Anatomical Landmarks for Euthanasia</i>) • Penetrating captive bolt devices with velocities ranging from at least 55–58 m/s are most effective <p>Secondary steps necessary to cause death:</p> <p>Secondary steps can ONLY be performed on an animal that is confirmed to be unconscious. Acceptable secondary steps include:</p> <ul style="list-style-type: none"> • bleeding out • pithing • rapid intravenous injection of a concentrated solution of potassium chloride or magnesium sulfate • a second shot (penetrating captive bolt or gunshot) when the above secondary methods are not available.

7.3 Confirming Loss of Consciousness and Death

Multiple indicators should be used when evaluating consciousness. Signs that an animal is unconscious include:

- no rhythmic breathing (breathing may be slow and erratic in an unconscious animal)
- animal does not blink when the surface of the eye is touched or show any eye movement
- no resistance is felt as the animal's jaw is opened wide; the jaw is relaxed
- no vocalization.

Animals euthanized by gunshot or penetrating captive bolt device should immediately be unconscious and collapse. Convulsions (i.e., uncoordinated kicking of the legs, body rigidity) following the application of any euthanasia method are not a sign of consciousness.

Humane methods of euthanasia are those that result in rapid loss of consciousness followed by death without the animal regaining consciousness (97). However, unlike loss of consciousness, death does not occur immediately: it is the result of respiratory and cardiac failure, which can take several minutes (98). Lack of movement is not an indicator of death. The following indicators should be used to confirm death in the 5 minutes after the application of the euthanasia method:

- lack of heartbeat (best evaluated with a stethoscope placed over the left lower chest area of the animal),
- lack of respiration, and
- no blinking when the eyeball is touched.

Carcasses must be disposed of in accordance with provincial regulations.

REQUIREMENTS

Cattle euthanized using gunshot or euthanasia drugs must be assessed to confirm that they are unconscious immediately after application. If an animal is not immediately unconscious, then a repeat application must be immediately delivered.

Cattle euthanized using a penetrating captive bolt must be assessed to confirm that they are unconscious immediately after application and before the secondary step to cause death is applied. If an animal is not immediately unconscious, then a repeat application must be immediately delivered.

Before moving or leaving the animal, death must be confirmed.



References

1. Adapted from: Dairy Cattle Welfare Council (2018) Principles of animal welfare. Available at: www.dwcouncil.org/node/4006. Accessed: December 3, 2022.
2. World Organisation for Animal Health (2019) Chapter 7.1 Introduction to the recommendations for animal welfare. In: *Terrestrial Animal Health Code, Section 7 – Animal Welfare*. Available at: www.woah.org/en/what-we-do/standards/codes-and-manuals/terrestrial-code-online-access/?id=169&L=1&htmlfile=chapitre_aw_introduction.htm. Accessed: December 17, 2022.
3. Government of Canada (2020) *Health of Animals Regulations C.R.C., c. 296*. Available at: www.laws-lois.justice.gc.ca/eng/regulations/c.r.c.,_c._296/page-10.html#h-548075. Accessed: September 28, 2020.
4. Dairy Cattle Code Scientific Committee (2020) Pain control for painful conditions and procedures. In: *Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues*. Lacombe AB: National Farm Animal Care Council.
5. American Veterinary Medical Association (AVMA) (2020) *AVMA Guidelines for the Euthanasia of Animals: 2020 Edition*. Schaumburg, IL: American Veterinary Medical Association. Available at: www.avma.org/sites/default/files/2020-02/Guidelines-on-Euthanasia-2020.pdf.
6. Dairy Cattle Code Scientific Committee (2020) Optimal management and design of indoor systems. In: *Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues*. Lacombe AB: National Farm Animal Care Council.
7. Rushen J. & de Passillé A.M. (2010) The importance of good stockmanship and its benefits for the animals. In: *Improving Animal Welfare: A Practical Approach*. Ed. T. Grandin. Cambridge MA: CAB International, pp. 50–63.
8. Grandin T. (2010) How to improve livestock handling and reduce stress. In: *Improving Animal Welfare: A Practical Approach*. Ed. T. Grandin. Cambridge MA: CAB International, pp. 64–87.
9. Dairy Cattle Code Scientific Committee (2020) Lameness and injuries. In: *Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
10. Van Os J. (2020–2021) Two heads are better than one: a starter guide to pairing calves. Available at: www.animalwelfare.cals.wisc.edu/calf_pairing. Accessed: December 17, 2022.
11. Costa J.H.C., von Keyserlingk M.A.G. & Weary D.M. (2016) Invited review: Effects of group housing of dairy calves on behavior, cognition, performance, and health. *Journal of Dairy Science* 99:2453–2467.
12. Miller-Cushon E.K. & DeVries T.J. (2016) Effect of social housing on the development of feeding behavior and social feeding preferences of dairy calves. *Journal of Dairy Science* 99(2):1406–1417.
13. Overvest M.A., Crossley R.E., Miller-Cushon E.K. & DeVries T.J. (2018) Social housing influences the behavior and feed intake of dairy calves during weaning. *Journal of Dairy Science* 101(9):8123–8134.
14. Bolt S.L., Boyland N.K., Mlynski D.T., James R. & Croft D.P. (2017) Pair housing of dairy calves and age at pairing: effects on weaning stress, health, production and social networks. *PLOS ONE* 12(1):e0166926.
15. Meagher R.K., Daros R.R., Costa J.H.C, von Keyserlingk M.A.G, Hötzel M.J. & Weary D.M (2015) Effects of degree and timing of social housing on reversal learning and response to novel objects in dairy calves. *PLOS ONE*. 10(8):e0132828.
16. Costa J.H.C., Meagher R.K., von Keyserlingk M.A.G. & Weary D.M. (2015) Early pair housing increases solid feed intake and weight gains in dairy calves. *Journal of Dairy Science* 98:381–386.



References (continued)

17. Veal Cattle Code of Practice Scientific Committee (2016) Comparison of the welfare implications of rearing veal calves in stall, tether, and group housing systems. In: *Code of Practice for the Care and Handling of Veal Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
18. Ede T., Weary D.M. & von Keyserlingk M.A.G. (2022) Calves are socially motivated. *JDS Communications* 3:44–48.
19. Keil N. M., Wiederkehr T.U., Friedli K. & Wechsler B. (2006) Effects of frequency and duration of outdoor exercise on the prevalence of hock lesions in tied Swiss dairy cows. *Preventive Veterinary Medicine* 74:142–153.
20. World Organisation for Animal Health (2017) Chapter 7.11 Animal welfare and dairy cattle production systems. In: *Terrestrial Animal Health Code*. Available at: www.woah.org/en/what-we-do/standards/codes-and-manuals/terrestrial-code-online-access/?id=169&L=1&htmlfile=chaptre_aw_dairy_cattle.htm. Accessed: December 17, 2022.
21. Proudfoot K. L., Jensen M. B., Weary D. M. & von Keyserlingk M.A.G. (2014) Dairy cows seek isolation at calving and when ill. *Journal of Dairy Science* 97:2731–2739.
22. Lorenz I., Earley B., Gilmore J., Hogan I., Kennedy E. & More S.J. (2011) Calf health from birth to weaning III. Housing and management of calf pneumonia. *Irish Veterinary Journal* 64(14).
23. Animal Health Ireland (n.d.) Control of pneumonia in dairy calves. Available at: www.animalhealthireland.ie/programmes/calfcare. Accessed: December 6, 2022.
24. Lorenz I., Fagan J. & More S.J. (2011) Calf health from birth to weaning II. Management of diarrhoea in pre-weaned calves. *Irish Veterinary Journal* 64(9).
25. Lactanet (2014) The Barn: A source of comfort. Practical guide to evaluating and improving comfort in the barn. Available at: www.lactanet.ca/en/practical-guide-to-evaluating-and-improving-comfort-in-the-barn. Accessed: November 8, 2022.
26. Leso L., Barbari M., Lopes M.A., Damasceno F.A., Galama P., Taraba J.L. & Kuipers K. (2019) Invited review: Compost-bedded pack barns for dairy cows. *Journal of Dairy Science* 103:1072–1099.
27. von Keyserlingk M.A.G., Barrientos A., Ito K., Galo E. & Weary D.M. (2012) Benchmarking cow comfort on North American freestall dairies: Lameness, leg injuries, lying time, facility design, and management for high-producing Holstein dairy cows. *Journal of Dairy Science* 95:7399–7408.
28. Veal Cattle Code of Practice Scientific Committee (2016) Flooring and bedding. In: *Code of Practice for the Care and Handling of Veal Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council. Available at: www.nfacc.ca/resources/codes-of-practice/veal-cattle/veal_cattle_SCreport_2016.pdf.
29. Rushen J., de Passillé A.M., von Keyserlingk M.A.G. & Weary D.M. (2008) Stockmanship and the interactions between people and cattle. In: *The Welfare of Cattle*. Vol. 5. Ed. C. Phillips. Dordrecht, NE: Springer, pp. 229–253.
30. Drackley J.K. (2016) The importance of BCS management to cow welfare, performance and fertility. *WCDS Advances in Dairy Technology* 28:195–206.
31. Roche S., Genore-Roche R. & Renaud D. (2021) *Perinatal mortality: A summary of current literature prepared for the dairy cattle Code Development Committee*. Lacombe, AB: National Farm Animal Care Council.
32. Esposito G. (2019) Ideal body condition score profile in dairy cows. *NDS Dynamics* 7(3). Available at: www.rumen.it/newsletter/481. Accessed: December 8, 2020.
33. Ferguson J. D. (1996) Implementation of a body condition scoring program in dairy herds. Centre for Animal Health and Productivity, University of Pennsylvania, School of Veterinary Medicine.



References (continued)

34. Endres M.I. & Espejo L.A. (2010) Feeding management and characteristics of rations for high-producing dairy cows in freestall herds. *Journal of Dairy Science* 93:822–829.
35. Bach A.N., Valls A.S. & Torrent T. (2008) Associations between nondietary factors and dairy herd performance. *Journal of Dairy Science* 91:3259–3267.
36. Rushen J., de Passillé A.M., von Keyserlingk M.A.G. & Weary D.M. (2008) Feeding and nutrition. In: *The Welfare of Cattle*. Vol. 5. Ed. C. Phillips. Dordrecht, NE: Springer, pp. 211–228.
37. DeVries T.J., von Keyserlingk M.A.G. & Beauchemin K.A. (2005) Frequency of feed delivery affects the behavior of lactating dairy cows. *Journal of Dairy Science* 88:3553–3562.
38. Sova A.D., LeBlanc S.J., McBride B.W. & DeVries T.J. (2013) Associations between herd-level feeding management practices, feed sorting, and milk production in freestall dairy farms. *Journal of Dairy Science* 96:4759–4770.
39. Yuan K. (2019) Why do we care about rumination? In: *Progressive Dairy*. Available at: www.progressivedairy.com/topics/feed-nutrition/why-do-we-care-about-rumination. Accessed: November 6, 2020.
40. Beauchemin K.A. (2018) Invited review: Current perspectives on eating and rumination activity in dairy cows. *Journal of Dairy Science* 101(6):4762–4784.
41. Dairy Cattle Code of Practice Scientists' Committee (2009) *Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
42. de Passillé A.M. & Rushen J. (2016) Using automated feeders to wean calves fed large amounts of milk according to their ability to eat solid feed. *Journal of Dairy Science* 99:3578–3583.
43. Neave H.W., Costa J.H.C., Benetton J.B., Weary D.M. & von Keyserlingk M.A.G. (2019) Individual characteristics in early life relate to variability in weaning age, feeding behavior, and weight gain of dairy calves automatically weaned based on solid feed intake. *Journal of Dairy Science* 102:10250–10265.
44. Veal Cattle Code of Practice Scientific Committee (2016) Management of milk feeding. In: *Code of Practice for the Care and Handling of Veal Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
45. Van Os J. (2020–2021) Feeding practices and reducing cross sucking. Available at: www.animalwelfare.cals.wisc.edu/wp-content/uploads/sites/243/2022/06/06-feeding.pdf. Accessed: November 8, 2022.
46. Montoro C., Miller-Cushon E.K., DeVries T.D. & Bach A. (2013) Effect of physical form of forage on performance, feeding behavior, and digestibility of Holstein calves. *Journal of Dairy Science* 96:1117–1124.
47. Bovine Alliance of Management & Nutrition (2017) A guide to feeding and weaning healthy and productive dairy calves. Available at: www.aphis.usda.gov/animal_health/nahms/dairy/downloads/bamn/BAMN17_GuideFeeding_1.pdf. Accessed: September 21, 2020.
48. Sweeney B.C., Rushen J., Weary D.M. & de Passillé A.M. (2010) Duration of weaning, starter intake, and weight gain of dairy calves fed large amounts of milk. *Journal of Dairy Science* 93:148–152.
49. Gelsinger S.L., Heinrichs A. J. & Jones C.M. (2016) A meta-analysis of the effects of preweaned calf nutrition and growth on first-lactation performance. *Journal of Dairy Science* 99:6206–6214.
50. PennState Extension (2016) Pasturing Dry Cows and Heifers. Available at: www.extension.psu.edu/pasturing-dry-cows-and-heifers. Accessed: September 18, 2020.
51. Tranel L. & Combs D. (n.d.) Feeding Dairy Cows on Quality Pasture. Available at: www.extension.iastate.edu/dairyteam/files/page/files/FeedingDairyCowsonQualityPasture.pdf. Accessed: September 18, 2020.



References (continued)

52. Wickramasinghe H.K.J.P., Kramer A.J. & Appuhamy J.A.D.R.N. (2018) Drinking water intake of newborn dairy calves and its effects on feed intake, growth performance, health status, and nutrient digestibility. *Journal of Dairy Science* 102:377–387.
53. Dairy Cattle Code Scientific Committee (2020) End-of-life management. In: *Code of Practice for the Care and Handling of Dairy Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
54. Canadian Veterinary Medical Association (2021) Position statement: Surgical procedures performed on animals. Available at: www.canadianveterinarians.net/policy-and-outreach/position-statements/statements/surgical-procedures-performed-on-animals. Accessed: December 17, 2022.
55. Canadian Veterinary Medical Association (2022) Position statement: Pain management in animals. Available at: www.canadianveterinarians.net/policy-and-outreach/position-statements/statements/pain-management-in-animals-previously-pain-control-in-animals-position-statement. Accessed: December 17, 2022.
56. Beef Code of Practice Scientists' Committee (2012) Painful procedures. In: *Code of Practice for the Care and Handling of Beef Cattle: Review of Scientific Research on Priority Issues*. Lacombe, AB: National Farm Animal Care Council.
57. American Veterinary Medical Association (2014) *Literature Review on the Welfare Implications of the Dehorning and Disbudding of Cattle*. Available at: www.avma.org/sites/default/files/resources/dehorning_cattle_bgnd.pdf. Accessed: June 4, 2020.
58. Canadian Veterinary Medical Association (2022) *Position statement: Horn management of cattle*. Available at: www.canadianveterinarians.net/policy-and-outreach/position-statements/statements/horn-management-of-cattle. Accessed: December 17, 2022.
59. Neely C.D., Thomson D.U., Kerr C.A. & Reinhardt C.D. (2014) Effects of three dehorning techniques on behavior and wound healing in feedlot cattle. *Journal of Animal Science* 92:2225–2229. doi:10.2527/jas2013-7424.
60. Canadian Veterinary Medical Association (2019) *Position statement: Castration of cattle, sheep, and goats*. Available at: www.canadianveterinarians.net/policy-and-outreach/position-statements/statements/castration-of-cattle-sheep-and-goats. Accessed: December 18, 2022.
61. Eicher S.D. & Dailey J.W. (2002) Indicators of acute pain and fly avoidance behaviors in Holstein calves following tail-docking. *Journal of Dairy Science* 85:2850–2858.
62. Kroll L.K., Grooms D.L., Siegford J.M., Schweihofer J.P., Daigle C.L., Metz K. & Ladoni M. (2014) Effects of tail docking on behavior of confined feedlot cattle. *Journal of Animal Science* 92:4701–4710.
63. Rushen J., de Passillé A.M., von Keyserlingk M.A.G. & Weary D.M. (2008) Stress and physiological indicators of animal welfare. In: *The Welfare of Cattle*. Vol. 5. Ed. C. Phillips. Dordrecht, NE: Springer, pp. 43–69.
64. Silanikove N., Merin U., Shapiro F. & Leitner G. (2013) Early mammary gland metabolic and immune responses during natural-like and forceful drying-off in high-yielding dairy cows. *Journal of Dairy Science* 96:6400–6411.
65. Dingwell R.T., Leslie K.E., Schukken Y.H., Sargeant J.M., Timms L.L., Duffield T.F., Keefe G.P., Kelton D.F., Lissemore K.D. & Conklin J. (2004) Association of cow and quarter-level factors at drying-off with new intramammary infections during the dry period. *Preventive Veterinary Medicine* 63(1–2):75–89.
66. Mastitis Network & Dairy Farmers of Canada (2020) Drying off cull dairy cattle at high production and in emergency situations. Available at: www.dairyresearch.ca/pdf/EN_Drying_off_PLIC_Aug62020FINAL.pdf. Accessed: November 6, 2020.



References (continued)

67. Jansen J. & Lam T.J. (2012) The role of communication in improving udder health. *Veterinary Clinics of North America: Food Animal Practice* 28:363–379.
68. Gulliksen S.M., Lie K.I. & Østerås O. (2009) Calf health monitoring in Norwegian dairy herds. *Journal of Dairy Science* 92(4):1660–1669.
69. Vasseur E., Rushen J., de Passillé A.M., Lefebvre D. & Pellerin D. (2010) An advisory tool to improve management practices affecting calf and heifer welfare on dairy farms. *Journal of Dairy Science* 93:4414–4426.
70. Lundborg G.K., Svensson E.C. & Oltenacu P.A. (2005) Herd-level risk factors for infectious diseases in Swedish dairy calves aged 0–90 days. *Preventive Veterinary Medicine* 68:123–143.
71. Lorenz I., Mee J.F., Earley B. & More S.J. (2011) Calf health from birth to weaning I: General aspects of disease prevention. *Irish Veterinary Journal* 64:10.
72. Canadian Veterinary Medical Association (2022) Position statement: Pest management. Available at: www.canadianveterinarians.net/policy-and-outreach/position-statements/statements/pest-management. Accessed: December 18, 2022.
73. Mee J.F. (2009) Bovine perinatology: Current understanding and future developments. In: *Animal Reproduction: New Research Developments*. Ed. L.T. Dahnof. New York, NY: Nova Science Publishers, pp. 67–106.
74. Schuenemann G.M., Bas S., Gordon E. & Workman J. (2011) Dairy calving management: Assessment of a Comprehensive Program for Dairy Personnel. *Journal of Dairy Science* 89–483.
75. Renaud D. (2019) Why do some calves thrive and others die? Risk factors impacting male and female dairy calf health. *Western Canadian Dairy Seminar Advances in Dairy Technology* 31:279–286.
76. Garthwaite B.D., Drackley J.K., McCoy G.C. & Jaster E.H. (1994) Whole milk and oral rehydration solution for calves with diarrhea of spontaneous origin. *Journal of Dairy Science* 77:835–843.
77. Based on data available through the National Dairy Study: Winder C.B., Bauman C.A., Duffield T.F., Barkema H.W., Keefe G.P., Dubuc J., Uehlinger F. & Kelton D.F. (2018) Canadian national dairy study: Heifer calf management. *Journal of Dairy Science* 101(11):10565–10579.
78. Renaud D.L., Kelton D.F., LeBlanc S.J., Haley D.B. & Duffield T.F. (2018) Calf management risk factors on dairy farms associated with male calf mortality on veal farms. *Journal of Dairy Science* 101:1785–1794.
79. Pyo J., Hare K., Pletts S., Inabu Y., Haines D., Sugino T., Guan L.L. & Steele M. (2020) Feeding colostrum or a 1:1 colostrum: milk mixture for 3 days postnatal increases small intestinal development and minimally influences plasma glucagon-like peptide-2 and serum insulin-like growth factor-1 concentrations in Holstein bull calves. *Journal of Dairy Science* 103(5):4236–4251.
80. Berge A.C.B., Besser T.E., Moore D.A. & Sischol W.M. (2008) Evaluation of the effects of oral colostrum supplementation during the first fourteen days on the health and performance of preweaned calves. *Journal of Dairy Science* 92:286–295.
81. Quigley J.D. & Wolfe T.M. (2003) Effects of spray-dried animal plasma in calf milk replacer on health and growth of dairy calves. *Journal of Dairy Science* 86:586–592.
82. Tyler J.W., Hancock D.D., Parish S.M., Rea D.E., Besser T.E., Sanders S.G. & Wilson L.K. (1996) Evaluation of 3 assays for failure of passive transfer in calves. *Journal of Veterinary Internal Medicine* 10(5):304–7.
83. McGuirk S.M. & Collins M. (2004) Managing the production, storage, and delivery of colostrum. *Veterinary Clinics Food Animal Practice* 20(3):593–603.
84. Windeyer M.C., Leslie K.E., Godden S.M., Hodgins D.C., Lissemore K.D. & LeBlanc S.J. (2014) Factors associated with morbidity, mortality, and growth of dairy heifer calves up to 3 months of age. *Preventive Veterinary Medicine* 113(2):231–240.
85. Rushen J., de Passillé A.M., von Keyserlingk M.A.G. & Weary D.M. (2008) Health, Disease, and Productivity. In: *The Welfare of Cattle*. Vol. 5. Ed. C. Phillips. Dordrecht, NE: Springer, pp. 15–42.



References (continued)

86. Sargeant J.M., Schukken Y.H & Leslie K.E. (1998) Ontario bulk milk somatic cell count reduction program: Progress and outlook. *Journal of Dairy Science* 81:1545–1554.
87. Ruegg, P. L. (2004) Pre-milking cow preparation – Secret methods of producing high quality milk. Proceedings of the regional meeting of the National Mastitis Council, pp. 34–40.
88. DeVries T.J. & von Keyserlingk M.A.G. (2005) Time of feed delivery affects the feeding and lying patterns of dairy cows. *Journal of Dairy Science* 88:625–631.
89. DeVries T.J., Dufour S. & Scholl D.T. (2010) Relationship between feeding strategy, lying behavior patterns, and incidence of intramammary infection in dairy cows. *Journal of Dairy Science* 93:1987–1997.
90. Cramer G. (n.d.) Addressing lameness: on the farm and at the industry level. Available at: www.dairyknow.umn.edu/topics/lameness/addressing-lameness-on-the-farm-and-at-the-industry-level. Accessed: May 7, 2020.
91. Canadian Food Inspection Agency (last updated 2020-11-10) *Health of Animals Regulations: Part XII: Transport of Animals – Regulatory Amendment Interpretive Guidance for Regulated Parties*. Available at: www.inspection.canada.ca/animal-health/terrestrial-animals/humane-transport/health-of-animals-regulations-part-xii/eng/1582126008181/1582126616914. Accessed: November 20, 2022.
92. Renaud D.L., Duffield T.F., LeBlanc S.J., Haley D.B & Kelton D.F. (2018) Clinical and metabolic indicators associated with early mortality at a milk-fed veal facility: A prospective case-control study. *Journal of Dairy Science* 101:2669–2678.
93. Wilson D. (2019) Predicting health outcomes and sale price of male dairy calves undergoing long distance transportation. MSc. Thesis, Applied Animal Biology. Vancouver: University of British Columbia.
94. Renaud D.L., Duffield T.F., LeBlanc S.J., Ferguson S., Haley D.B & Kelton D.F. (2018) Risk factors associated with mortality at a milk-fed veal calf facility: A prospective cohort study. *Journal of Dairy Science* 101:2659–2668.
95. Winder C.B., Kelton D.F. & Duffield T.F. (2016) Mortality risk factors for calves entering a multi-location white veal farm in Ontario, Canada. *Journal of Dairy Science* 99:10174–10181.
96. Grandin, T. Livestock handling systems, cattle corrals, stockyards, and races. Available at: www.grandin.com/design/design.html. Accessed: October 8, 2020.
97. Canadian Veterinary Medical Association (CVMA) (2021) *Position statement: Euthanasia*. Available at: www.canadianveterinarians.net/documents/euthanasia. Accessed: December 19, 2022.
98. Ontario Association of Bovine Practitioners (2019) Considerations for developing a down cattle protocol. Available at: www.oabp.ca/images/news/Considerations-for-developing-a-down-cattle-protocol-November-6-2019.pdf. Accessed: December 19, 2022.
99. Shearer J.K. & Ramirez A. (last updated January 28, 2013) Procedures for Humane Euthanasia – Euthanasia of Sick, Injured and/or Debilitated Livestock. Available at: www.vetmed.iastate.edu/sites/default/files/vdpam/Extension/Dairy/Programs/Humane%20Euthanasia/Download%20Files/EuthanasiaBrochure20130128.pdf. Accessed: May 26, 2020.
100. Humane Slaughter Association (2018) Humane Dispatch and Disposal of Infant Calves: Technical Note No. 2. Available at: www.hsa.org.uk/downloads/technical-notes/tn2-humane-dispatch-of-infant-calves.pdf. Accessed: May 5, 2020.
101. Woods J., Shearer J.K. & Hill J. (2010) Recommended On-Farm Euthanasia Practices. In: *Improving Animal Welfare: A Practical Approach*. Ed. Temple Grandin. Cambridge, UK: CAB International, pp. 186–213.



Sample Cattle Welfare Policy

[Your Farm/Company]

Employee Animal Care Code of Conduct

Our commitment to our animals

[Our company/farm] is committed to responsible farm animal care and handling. That means animals in our care deserve to be healthy, safe and well cared for.

Our commitment to our customers

Working with animals is important work that we take seriously. We are proud of the work that we do, and we strictly enforce responsible farm animal care and handling among employees and service providers at our facility.

Every person who handles or comes into contact with an animal is required to support our core objective of responsible farm animal care and handling. The demonstration of that support is through the review and signing of this Code of Conduct agreement on a [quarterly/annual] basis.

Our commitment to our employees

Your job is valuable and important to our animals, and our business. When you report an incident involving possible mistreatment, illness or injury involving one of our animals, we will take it seriously. We will document your concern. We will follow up to resolve the animal's situation, and/or provide additional training among employees.

Our employees' commitment to us

Every one of our employees is required to handle and treat animals with respect and in accordance with [farm/company] policies and rules as well as the federal, provincial, and municipal regulations under which we operate.

Any employee who is responsible for, observes or receives any information that alleges an animal on our property or in our care is being mistreated, mishandled or treated or handled in a way that is contrary to our animal care policy/guidelines must report that information to [NAME OF POINT PERSON] immediately so that the situation can be corrected. [PROVIDE CONTACT INFO].

Failure to adhere to this agreement is cause for dismissal. [Farm/company] reserves the right to refer animal-abusers to law enforcement for prosecution.

I _____ understand and acknowledge that willful neglect, mishandling or abuse of animals by any [name of company] employee or witnessing it and not reporting it is subject to discipline including immediate termination of employment, and that offenders may also be subject to prosecution under applicable laws.

Signature of employee

Date

Name (Please Print)

Signature of Employer

Date

Name

Title

Important Note: Seek advice from your legal counsel and human resources department if appropriate to ensure any agreement meets relevant labour laws and union contracts.

Used with permission from Farm & Food Care Ontario.


www.farmfoodcareon.org/wp-content/uploads/2016/06/Animal-Care-Code-of-Conduct-2016.pdf



Body Condition Scoring Charts

LABELLED ILLUSTRATION OF A DAIRY COW

Illustration of a Dairy Cow with a BCS score of 3.



BODY CONDITION SCORES FOR DAIRY COWS
Overview of all the body condition scores for Dairy Cows

BCS 1:

SHORT RIBS:

- Ends sharp to touch
- Loin prominent, shelf-like appearance
- Obvious scalloping over top and ends

BACKBONE:

- Vertebrae prominent in chine, loin and rump area
- Individual bones easily visible

HOOK AND PIN BONES:


- Sharply defined, very angular in appearance
- No discernable fat pad

THURL (area over pelvis):

- Severe “V shaped” depression without fat cover

TAIL HEAD:

- Sunken and hollow on either side of tail head with obvious folds of skin
- Ligaments connecting pin bones to spine are sharply defined
- Vulva prominent.



BCS 2

SHORT RIBS:

- Ends not as prominent as BCS 1, but can be felt
- Edges easily felt, with slight fat cover, and slightly more rounded appearance
- Overhanging shelf effect less apparent

BACKBONE:

- Vertebrae in chine, loin and rump area, less visually distinct
- Easily feel individual vertebrae

HOOK AND PIN BONES:


- Bones still prominent, angular
- No fat pad palpable

THURL (area over pelvis):

- Less severe “V shaped” depression
- Little tissue cover

TAIL HEAD:

- Both sides of the tail head are sunken and hollow
- Sharply defined ligaments connecting pin bones to spine





Body Condition Scoring Charts (continued)

BODY CONDITION SCORES FOR DAIRY COWS

Overview of all the body condition scores for Dairy Cows

BCS 3

SHORT RIBS:

- Ends can be felt with moderate pressure
- Ribs appear smooth without noticeable scalloping
- Overhanging shelf effect much less apparent

BACKBONE:

- Vertebrae in chine, loin and rump area appear rounded
- Backbone visible, but individual vertebrae not distinct

HOOK AND PIN BONES:


- Visible, but smooth, with rounded appearance
- Fat pad palpable

THURL (area over pelvis):

- Forms “U shaped” depression

TAIL HEAD:

- Both sides of tail head somewhat hollow, but skin folds not distinct
- Ligaments connecting pin bones to spine are rounded in appearance



BCS 4

SHORT RIBS:

- Individual rib ends not visible, only felt with firm pressure
- Overhanging shelf effect slight, barely visible

BACKBONE:

- Vertebrae in chine rounded, smooth
- Loin and rump areas appear flat

HOOK AND PIN BONES:


- Rounded, with obvious fat covering

THURL (area over pelvis):

- Area between hooks and pins almost flat
- Pelvic bone only felt with firm pressure

TAIL HEAD:

- Sides of tail head not hollow, no skin folds
- Some fat deposit palpable



BCS 5

SHORT RIBS:

- Ends can't be seen or felt
- No overhanging shelf effect

BACKBONE:

- Vertebrae in chine, loin and rump not visible
- Difficult to feel individual vertebrae

HOOK AND PIN BONES:


- Very round, buried (almost disappearing) in fat tissue

THURL (area over pelvis):

- Appears flat
- Filled in between the hooks and pins

TAIL HEAD:

- Hollow filled in
- Areas on both sides of tail head buried in fat tissue



Source: Adapted from *What's the Score? Body Condition Scoring for Livestock* CD-ROM, Agdex CD 400/40-1. Used with permission from Alberta Agriculture and Rural Development. Available at: www.agric.gov.ab.ca/app08/ppsropintheweb?PubID=100031



Body Condition Scoring Charts (continued)

Quarter Point Body Condition Scoring Chart

This is a 14-point body condition scoring chart (with scores ranging from <2 to 5 in 0.25 increments). Scores are split between thinner (BCS of 3.0 and lower; top table) and fatter cows (BCS of 3.25 and higher; bottom table). Gray cells depict what has changed between two scores. Pelvic area is for either a V-shaped (thinner) or U-shaped (fatter) cow. Start by determining if the pelvic area is V-shaped or U-shaped, and then work from left to right on the relevant table (V-shaped is the top table, U-shaped is the bottom table).

BCS	3.0	2.75	2.5	2.25	2.0	<2.0
Pelvic Area	V	V	V	V	V	V
Hook Bones	rounded	angular	angular	angular	angular	angular
Pin Bones	padded	padded	angular, fat palpable	angular, no fat palpable	angular, no fat palpable	angular, no fat palpable
Short Ribs	corrugations between ribs not visible	corrugations between ribs not visible	corrugations between ribs not visible	corrugations visible ½ way between tips and spine	corrugations visible ¾ way between tips and spine	corrugations visible all the way up to the spine

BCS	3.25	3.5	3.75	4.0	4.25	4.5	4.75	5.0
Pelvic Area	U	U	U	U	U	U	U	U
Tailhead Ligament	visible	barely visible	not visible	not visible	not visible	not visible	not visible	not visible
Sacral Ligament	visible	visible	barely visible	not visible	not visible	not visible	not visible	not visible
Thurl	not flat	not flat	not flat	not flat	flat	flat	flat	flat
Tips of Short Ribs	visible	visible	visible	visible	barely visible	barely/not visible	barely/not visible	barely/not visible
Pin Bones	visible	visible	visible	visible	visible	buried	buried	buried
Hook Bones	visible	visible	visible	visible	visible	visible	barely visible	not visible

Adapted from Vasseur E., Gibbons J., Rushen J. & de Passillé A.M. (2013) Development and implementation of a training program to ensure high repeatability of body condition scoring of dairy cows. Journal of Dairy Science 96(7):4725–4737.



Cow Cleanliness Scoring

Cow Cleanliness Assessment



Réseau canadien de recherche sur la mammité bovine
Canadian Bovine Mastitis Research Network



The cleanliness of cows has a significant impact on udder health and more particularly on the rate of environmental mastitis. Maintaining a clean udder and legs helps reducing the spread of environmental pathogens to the teat canal. Depending on what part of the cow is soiled, it is possible to determine what areas of the barn have an inadequate level of cleanliness, therefore appropriate corrective action can be taken.

	1	2	3	4
Udder				
Hind legs				
Flanks and hips				

Adapted from fact-sheets produced by the Faculty of Veterinary Medicine, University of Wisconsin, and Pfizer Animal Health. Most photos courtesy of Pfizer Animal Health.

Photo credits: *«Less Mastitis, Better Milk»* (Pierre Lévesque, 2004), distributed by Hoard's Dairyman (www.hoards.com).

Udder cleanliness

(back and sides)

is an indicator of the cleanliness of stalls and bedding.

(Examine right before milking)

If the standard is not met, check:

- Cleanliness of stalls
- Amount of bedding
- Need to shave/singe udder hair
- Manure consistency

Hind legs cleanliness

is an indicator of the cleanliness of alleyways and the length of tie stalls.

If the standard is not met, check:

- Cleanliness of alleyways and exterior areas
- Cleanliness of holding area
- Dimension of stalls
- Manure consistency

Flanks and hips cleanliness

is an indicator of the cleanliness of stalls and bedding.

If the standard is not met, check:

- Cleanliness of stalls
- Amount of bedding
- Manure consistency













This document may be reproduced in its entirety only, for educational purposes, without obtaining permission. If credit is given to CBMQRN.

01/09/10

Source: Canadian Bovine Mastitis and Milk Quality Research Network (CBMQRN)



Calf Health Scoring Chart and Criteria (continued)

Calf Health Scoring Criteria			
0	1	2	3
Nose Score – nasal discharge			
Normal serous discharge	Small amount of unilateral cloudy discharge	Bilateral cloudy or excessive mucus discharge	Copious bilateral mucopurulent discharge
			
Eye Score - discharge			
Normal – no discharge	Small amount of ocular discharge	Moderate amount of bilateral discharge	Heavy ocular discharge
			
Ear Score			
Normal	Ear flick or head shake	Slight unilateral droop	Head tilt or bilateral droop
			
Cough Score			
None	Induce single cough	Induced repeated or occasional spontaneous coughs	Repeated spontaneous coughs
Temperature Score			
100-100.9°F or 37.8-38.2°C	101-101.9°F or 38.3-38.8°C	102-102.9°F or 38.9-39.4°C	>103°F or >39.4°C
Fecal Score			
Normal	Semi-formed, pasty	Loose, but stays on top of bedding	Water, sifts through bedding
Navel Score			
Normal	Slightly enlarged, not warm or painful	Slightly enlarged, with slight pain, heat or moisture	Enlarged with pain, heat or malodorous discharge
Joint Score			
Normal	Slight swelling, not warm or painful	Swelling with pain or heat	Swelling with severe pain, heat or dislocation

Appendix D is used with permission from University of Wisconsin-Madison, School of Veterinary Medicine.



Lameness Scoring System for Dairy Cows

Table E.1 is for scoring cows in free stall barns; see next page for scoring cows in tie stall barns.

Table E.1 Gait Scoring Cows in Free Stall Barns

Score	Description	Behavioural Criteria
1 Sound	Smooth and fluid movement	<ul style="list-style-type: none"> • Flat back when standing and walking • All legs bear weight equally • Joints flex freely • Head carriage remains steady as the animal moves
2	Imperfect locomotion but ability to move freely not diminished	<ul style="list-style-type: none"> • Flat or mildly arched back when standing and walking • All legs bear weight equally • Joints slightly stiff • Head carriage remains steady
3	Capable of locomotion but ability to move freely is compromised	<ul style="list-style-type: none"> • Flat or mildly arched back when standing; possible arch when walking • Slight limp can be discerned in one or more limbs • Joints show signs of stiffness but do not impede freedom of movement • Head carriage remains steady
4	Ability to move freely is obviously diminished	<ul style="list-style-type: none"> • Obvious arched back when standing and walking • Reluctant to bear weight on at least one limb but still uses that limb in locomotion • Strides are hesitant and deliberate and joints are stiff • Head bobs slightly as animal moves in accordance with the sore hoof making contact with the ground
5 Severely Lame	Ability to move is severely restricted Must be vigorously encouraged to stand and/or move	<ul style="list-style-type: none"> • Extreme arched back when standing and walking • Inability to bear weight on one or more limbs • Obvious joint stiffness characterized by lack of joint flexion with very hesitant and deliberate strides • One or more strides definitely shortened • Head obviously bobs as sore hoof makes contact with the ground
Source: University of British Columbia Animal Welfare Program		

Table E.1 is adapted from Flower F.C. & Weary D.M. (2006) *Effect of hoof pathologies on subjective assessments of dairy cow gait*. Journal of Dairy Science 89:139–146.



Lameness Scoring System for Dairy Cows (continued)

Table E.2 Lameness Scoring Cows in Tie Stall Barns

Behaviour Indicator	Description
Standing Post (voluntary movements)	
EDGE	Placement of one or more hooves on the edge of the stall while standing stationary. Standing on the edge of a step when stationary, typically to relieve pressure on one part of the claw (Figures 1 and 2). This does not refer to when both hind hooves are in the gutter or when cow briefly places her hoof on the edge during a movement/step.
WEIGHT SHIFT	Regular, repeated shifting of weight from one hoof to another. Repeated shifting is defined as lifting each hind hoof at least twice off the ground (L-R-L-R or vice versa). The hoof must be lifted and returned to the same location and does not include stepping forward or backward.
REST (UNEVEN WEIGHT)	Repeated resting of one foot more than the other as indicated by the cow raising a part or the entire hoof off the ground and placing it back in the same spot. This does NOT include raising of the hoof to lick or during kicking or taking a step forward or back.
Cow moved from side to side	
UNEVEN MOVEMENT	Uneven weight bearing between hooves when the cow is encouraged to move from side to side. This is demonstrated by greater rapid movement of one hoof relative to the other, or by an evident reluctance to bear weight on a particular foot.



Figure 1: Example of EDGE




Figure 2: Example of EDGE

Source: Gibbons J., Haley D.B, Higginson Cutler J., Nash C., Zaffino J., Pellerin D., Adam S., Fournier A., de Passillé A.M., Rushen J. & Vasseur E. (Submitted) Technical Note: Reliability and Validity of a Method to Measure Lameness Prevalence of Cows in Tie-stalls. Journal of Dairy Science.



Transport Decision Tree




IMPORTANT: Lactating cows must reach their final destination (or a suitable place for milking) before becoming engorged. Cows with mammary engorgement will be considered compromised or unfit. Options include milking to prevent engorgement, drying off, or shipping when their milk production has decreased.

For more information, visit www.inspection.gc.ca/humane



Sample Euthanasia Decision Tree





Anatomical Landmarks for Euthanasia




Figure H.1

**Not Between the Eyes! -
but above the eyes as
illustrated.**




Figure H.2

Images reprinted with permission: J.K. Shearer and A. Ramirez, College of Veterinary Medicine, Iowa State University www.vetmed.iastate.edu/HumaneEuthanasia (2013).

Proper positioning of the firearm or penetrating captive bolt is necessary to achieve the desired results.

Figure H.1: In mature cattle, the correct target is in the middle of the forehead at the intersection of 2 imaginary lines drawn from the outside corner of each eye to the opposite horn or equivalent site in hornless animals. NOT between the eyes or at the poll.

Figure H.2: The firearm or captive bolt should be angled so the projectile follows the angle of the neck or spine. Ensure the aim is perpendicular to the skull but tilted slightly to direct the shot to the lower brain.



Anatomical Landmarks for Euthanasia (continued)



Figure H.3

Image source: Code of Practice for the Care and Handling of Veal Cattle (2017) Lacombe AB: National Farm Animal Care Council.

Figure H.3: In young calves, the correct target is in the middle of the forehead at the intersection of 2 imaginary lines drawn from the outside corner of each eye to the opposite horn or equivalent site in hornless animals. Because the forebrain of calves is underdeveloped (compared to older cattle), it is beneficial to direct the projectile towards the base of the skull.

When using a firearm: For all weight and age classes, the firearm must never be held in direct contact with the head.

When using a penetrating captive bolt gun: For all weights and ages, the device must be held in contact with the head using proper landmarks (figures H.1, H.2, H.3). Restraint may be necessary to ensure proper application of the captive bolt gun. A rope halter is typically sufficient to restrain the head. Alternatively, sedation may be used (in consultation with the herd veterinarian) as a means of humane restraint.



Secondary Steps to Cause Death

Bleeding out (exsanguination)

Bleeding out can only be performed on an unconscious animal using a very sharp knife with a rigid blade at least 15 cm (6 in) in length.¹



Figure I.1

Figure I.1. Bleeding out of an unconscious animal (previously stunned with captive bolt device or gunshot):

Insert a sharp, single-sided blade, at least 15 cm (6 in) long into the neck below the neck bones and behind the jaw. Draw the blade forward to sever the major blood vessels (jugular vein and carotid artery) of the neck and the windpipe (trachea). Blood should begin to flow freely, and death occurs within minutes.

Pithing

Pithing is the process of mechanically destroying the brain of an unconscious animal to prevent a return to consciousness.¹ Pithing is performed by inserting a rod or cane (approximately 1 m [3 ft] long x 5–10 mm [0.2–0.4 in] in diameter) through the hole in the skull created by the penetrating captive bolt device (Figure I.2).² Pithing rods are commercially available (e.g., www.pithingrods.com). The carcass is no longer safe for consumption due to possible contamination.¹ Producers are also advised to confirm that pithing will not affect dead stock removal.




Figure I.2

Figure I.2. Pithing of a previously stunned animal. *The line perpendicular to the animal's forehead shows the location for application of the captive bolt device. The curved pithing rod (shown only partially inserted) is inserted into the hole in the skull created by stunning with a penetrating captive bolt device. The rod is then manipulated, moving back and forth in approximately 5 cm (2 in) increments destroying the brain tissue and travelling through the brain to the top of the spinal cord, ensuring death.*



Secondary Steps to Cause Death (continued)

Rapid Intravenous Injection

A concentrated solution of potassium chloride or magnesium sulfate can be used. These drugs must only be administered to an unconscious animal. Administration of a saturated potassium chloride solution at a rate of 75–100 mg/kg body weight induces cardiac arrest. Example: an average mature cow at 636 kg (1,400 lb) would need 3–4 syringes of 1tbsp (20g) potassium chloride mixed in 60 ml of water each.³ Any available vein may be used; the handler should be positioned out of the reach of legs which may injure the operator during periods of involuntary movement.⁴ Once the needle is in the vein, the injection should be delivered by rapid intravenous injection.⁴

Biosecurity Considerations:

Disposal of blood is an important consideration when bleeding out is used as a secondary step for humane euthanasia; strategies include:

- use sawdust, wood shavings, straw, or other absorbent material to contain the blood and dispose of the material as required by provincial regulations for deadstock disposal
- if infectious disease is suspected, non-porous surfaces (e.g., floors, walls, equipment) should be cleaned and disinfected after bleeding out
- if infectious disease is suspected and the surface is soil, gravel, sand, or similar material: remove the top 20 cm (7.9 in) of material where the blood spill occurred and dispose of it as required by provincial regulations for deadstock disposal. Allow the area to dry before allowing contact with other animals.

¹ American Veterinary Medical Association (2020) *AVMA Guidelines for the Euthanasia of Animals: 2020 Edition*. Available at: www.avma.org/KB/Policies/Documents/euthanasia.pdf.

² Appelt M. & Sperry J. (2007) Stunning and killing cattle humanely and reliably in emergency situations – A comparison between a stunning-only and a stunning and pithing protocol. *Canadian Veterinary Journal* 48:529–534.

³ American Association of Bovine Practitioners (2019) *Guidelines for the Humane Euthanasia of Cattle*. Available at: www.aabp.org/Resources/AABP_Guidelines/EUTHANASIA-2019.pdf.

⁴ Shearer J.K. & Ramirez A. (2013) *Procedures for Humane Euthanasia – Euthanasia of Sick, Injured and/or Debilitated Livestock*. Available at: www.vetmed.iastate.edu/sites/default/files/vdpam/Extension/Dairy/Programs/Humane%20Euthanasia/Download%20Files/EuthanasiaBrochure20130128.pdf.



Resources for Further Information

PRODUCER MENTAL HEALTH SUPPORTS

- Canadian Mental Health Association www.cmha.ca
- The Do More Agriculture Foundation www.domore.ag
- Au coeur des familles agricoles (ACFA). La maison ACFA www.acfareseaux.qc.ca/fr/maison-acfa

HOUSING

Heifers and Cows

- Valacta (2014) The Barn: A source of comfort. Practical guide to evaluating and improving comfort in the barn. Available at: www.valacta.com/en-CA/gpc/_media/Document/guide-confort-ang-v3-version-finale-2015-02-02.pdf

Calves

- University of Wisconsin-Madison (2021) Two heads are better than one: starter guide to pairing dairy calves. Available at: www.animalwelfare.cals.wisc.edu/calf_pairing Accessed: September 16, 2021
- Cantor M.C, Neave H.W. & Costa J.H.C. (2020) Effectively raising pair-housed calves: Common questions from transitioning farmers. Progressive Dairy. Available at: www.progressivedairy.com/topics/calves-heifers/effectively-raising-pair-housed-calves-common-questions-from-transitioning-farmers Accessed: August 31, 2021

Ventilation

- Calfcare.ca (n.d.) Ventilation. Available at: www.calfcare.ca/housing/ventilation
- House H. (n.d.) Positive pressure air tube ventilation for calf housing. Ontario Ministry of Food and Rural Affairs, Agdex #420/721. Available at: www.ontario.ca/page/positive-pressure-air-tube-ventilation-calf-housing
- Nordlund K. Practical considerations for ventilating calf barns in winter. American Association of Bovine Practitioners, 40th Annual Conference, September 18, 2007. Vancouver, BC, Canada. Available at: www.fyi.extension.wisc.edu/heifermgmt/files/2015/02/Practical_ventilating_calf_barns.pdf

Emergency Planning

- Government of British Columbia (n.d.) *Emergency Planning Workbook for B.C. Dairy Producers*. Available at: www2.gov.bc.ca/assets/gov/farming-natural-resources-and-industry/agriculture-and-seafood/farm-management/emergency-management/dairy_emergency_management_guide.pdf. Accessed September 17, 2021
- Ontario Ministry of Agriculture, Food and Rural Affairs (n.d.) *Barn fires – A Concern for Ontario Farmers: Questions and answers to barn fires and fires in farm structures*. Available at: www.omafra.gov.on.ca/english/engineer/facts/barn_fire.htm
- Ontario Ministry of Agriculture, Food and Rural Affairs (2011) *Reducing the Risk of Fire on Your Farm*. Publication 837. Available at: www.omafra.gov.on.ca/english/engineer/barnfire/toc.pdf
- Ontario Ministry of Agriculture, Food and Rural Affairs (2016) *Electrical Systems in Barns*. Available at: www.farmfoodcareon.org/wp-content/uploads/2016/04/Electrical-Systems-in-Barns.pdf
- The Alberta Environmental Farm Plan Company (2008) *Rural Emergency Plan*. Available at: www.ruralemergencyplan.com



Resources for Further Information (continued)

FEED AND WATER

Assessing Rumen Fill

- Elanco Animal Health (2018) Rumen fill scoring. Available at: www.cmapspublic3.ihmc.us/rid=1WBBHH2N3-1W2284N-1KVM/rumen%20fill%20scoring.pdf
- Agriculture and Horticulture Development Board (n.d.) Rumen fill score card. Available at: www.projectblue.blob.core.windows.net/media/Default/Dairy/Publications/RumenScorecard2881_200702_WEB.pdf

Calf Feeding and Weaning

- Steele M. & Lallemand Animal Nutrition (2019) Healthy gut, healthy calf, productive future. Available at: www.youtube.com/watch?v=zFftJ6fw55w
- Lactanet (2021) Growth chart by breed. Available at: www.lactanet.ca/en/growth-chart-by-breed
- Bovine Alliance of Management and Nutrition (2008) Guide to calf milk replacers: types, use and quality. Available at: www.aphis.usda.gov/animal_health/nahms/dairy/downloads/bamn/BAMN08_GuideMilkRepl.pdf
- Quigley J. (2007) *Added CMR feeding in cold weather*. Calf note #121. Available at: www.calfnotes.com/pdffiles/CN121.pdf

Water

- Olkowski A.A. (2009) Livestock Water Quality: A Field Guide for Cattle, Horses, Poultry and Swine. Available at: www5.agr.gc.ca/resources/prod/doc/terr/pdf/lwq_guide_e.pdf
- Ontario Ministry of Agriculture, Food and Rural Affairs (2019) Water requirements of livestock. Available at: www.omafra.gov.on.ca/english/engineer/facts/07-023.htm

HUSBANDRY PRACTICES

Handling and Restraint

- Farm and Food Care Ontario (n.d.) Handling cattle: Work smarter not harder. Available at: www.farmfoodcareon.org/wp-content/uploads/2016/04/CattleHandlingYouth.pdf
- Grandin T. (Revised 2015) Understanding Flight Zone and Point of Balance for Low Stress Handling of Cattle, Sheep, and Pigs. Available at: www.grandin.com/behaviour/principles/flight_zone.html
- Grandin T. (Revised 2017) The Principles of Low Stress Restraint Cattle and Pigs. Available at: www.grandin.com/restrain/rest princ.html
- Gill R. & Machen R. (n.d.) Cattle handling pointers. Texas A&M AgriLife Extension. Available at: www.effectivestockmanship.com/PDFs/Cattle-Handling-Pointers.pdf
- University of Minnesota. Stockmanship. Available at: www.dairyknow.umn.edu/topics/stockmanship

Dry-Off Management

- Mastitis Network and Dairy Farmers of Canada (2020) Drying off cull dairy cattle at high production and in emergency situations. Available at: www.dairyresearch.ca/pdf/EN_Drying_off_PLC_Aug62020FINAL.pdf



Resources for Further Information (continued)

HEALTH MANAGEMENT

General

- Farmers Assuring Responsible Management (n.d.) Herd health plan template. Available at: www.nationaldairyfarm.com/wp-content/uploads/2018/11/Blank-Herd-Health-Plan.pdf Accessed: January 14, 2020.
- Dairy Farmers of Canada. Animal care and health factsheets (various topics) Available at: www.dairyfarmers.ca/proaction/resources/animal-care Accessed: September 22, 2021.

Lameness

- Dairy Farmers of Canada (2017) Lameness information document. Available at: www.dairyresearch.ca/pdf/EN_PLC_boiterie_fin.pdf
- University of Minnesota. Lameness (website – online articles, presentations, and videos). Available at: www.dairyknow.umn.edu/topics/lameness
- University of Wisconsin-Madison. Lifestep Lameness Module. Available at: www.thedairylandinitiative.vetmed.wisc.edu/home/lifestep-lameness-module

Respiratory Health

- Nordlund K. (n.d.) Housing factors to optimize respiratory health of calves in naturally ventilated calf barns in winter. Available at: www.vetmed.wisc.edu/dms/fapm/fapmtools/8calf/Calf_Barn_Ventilation_Text.pdf

Calf Health

- Charlton, S.J. (2009) Calf Rearing Guide. Copies can be ordered through Context Bookshop: www.contextbookshop.com/books/calf-rearing-guide-practical-easy-to-use
- Grober Nutrition. (2020) What three things should you consider for your colostrum program? Available at: www.grobernutrition.com/grofacts/what-three-things-should-you-consider-for-your-colostrum-program
- Quigley J. (2009) Prolonged colostrum feeding and calf health. Calf note #138. Available at: www.calfnotes.com/pdffiles/CN138.pdf

Calving

- Animal Health Ireland (2011) Calving and Care of the Newborn Calf. Available at: www.yumpu.com/en/document/view/5367781/calving-and-care-of-the-newborn-calf-animal-health-ireland

Down Cattle

- Ontario Association of Bovine Practitioners (2019) Considerations for developing a down cattle protocol. Available at: www.oabp.ca/images/news/Considerations-for-developing-a-down-cattle-protocol-November-6-2019.pdf
- Ontario Association of Bovine Practitioners (2019) Quick guide: down cattle management. Available at: www.oabp.ca/images/news/Down-Cattle-Quick-Guide-August-12-2019.pdf

Federally Reportable Diseases

- Canadian Food Inspection Agency (last modified 2020-08-07) Terrestrial animal diseases. Available at: www.inspection.gc.ca/animals/terrestrial-animals/diseases/eng/1300388388234/1300388449143



Resources for Further Information (continued)

TRANSPORTATION

- Dairy Farmers of Canada. Animal Care: Cull Cow Resources. Available at: www.dairyfarmers.ca/proaction/resources/animal-care
- Grandin T. (2008) Engineering and design of holding yards, loading ramps and handling facilities for land and sea transport of cattle. *Veterinaria Italiana* 44(1):235-245. Available at: www.researchgate.net/publication/43202698_Engineering_and_design_of_holding_yards_loading_ramps_and_handling_facilities_for_land_and_sea_transport_of_livestock
- Canadian Food Inspection Agency (last updated 2020-11-10) *Health of Animals Regulations: Part XII: Transport of Animals-Regulatory Amendment Interpretive Guidance for Regulated Parties*. Available at: www.inspection.gc.ca/animal-health/humane-transport/health-of-animals-regulations-part-xii/eng/1582126008181/1582126616914
- Government of Canada (2020) Health of Animals Regulations C.R.C., c. 296. Available at: www.laws-lois.justice.gc.ca/eng/regulations/c.r.c.,_c._296/page-15.html#h-548075
- Ontario Ministry of Agriculture and Food, and Ministry of Rural Affairs (n.d.) Ontario Livestock Manifest. Available at: www.farmfoodcareon.org/wp-content/uploads/2016/04/OntarioCattleTransportationManifest.pdf
- Certified Livestock Transport. Available at: www.livestocktransport.ca/en



Participants

Code Development Committee Members

Role	Representative	Organization
Producer	David Wiens (Chair)	Dairy Farmers of Canada
	Dave Taylor	Dairy Farmers of Canada
	Gerrit Damsteegt	Dairy Farmers of Canada
	Pascal Leduc Yvan Bastien (2019–2022)	Dairy Farmers of Canada
	Steve Runnalls	Dairy Farmers of Canada
Veterinarian	Kelly Barratt DVM	Canadian Veterinary Medical Association
National animal welfare association	Jeff Rushen PhD	Humane Canada
Provincial animal protection enforcement authority	Mike Draper	Ontario Ministry of the Solicitor General
Provincial government representative with responsibilities in animal welfare	Jane Pritchard DVM, MVSc	British Columbia Ministry of Agriculture and Food (retired)
Processor	Marie Ly	Dairy Processors Association of Canada
Research/Academic	Elsa Vasseur PhD	Scientific Committee Co-Chair
	Trevor DeVries PhD	Scientific Committee Co-Chair
Program implementation expert	Chantal Fleury agr.	Les Producteurs de lait du Québec
	Guy Séguin P.Eng. Maria Leal (2019–2021)	Dairy Farmers of Ontario
Technical expert	Steve Adam agr.	Lactanet
Federal government	Jean Lambert MA Lucille McFadden (2019–2021)	Agriculture and Agri-Food Canada
Allied industry	Kirk Jackson	Canadian Cattle Association
	Brian Keunen MSc	Canadian Veal Association
Industry liaison (Ex-Officio)	Nicole Sillett	Dairy Farmers of Canada

Scientific Committee Members

Name	Named by
Elsa Vasseur PhD (Co-Chair)	International Society for Applied Ethology
Trevor DeVries PhD (Co-Chair)	International Society for Applied Ethology
Charlotte Winder DVM, DVSc	Canadian Veterinary Medical Association
Daniel M. Weary DPhil	Canadian Society of Animal Science
Todd Duffield DVM, DVSc	Canadian Veterinary Medical Association

Participants are defined as per NFACC's Guiding Principles for Codes of Practice.

The Code Development Committee would like to thank Olivier Beaulieu-Charbonneau and Sandra Hamilton who also served on the committee for a time and whose perspectives were valued. We also thank Jackie Crichton, Kendra Keels, Sophie Neveux, and Warren Skippon who all contributed to this process as Standing Observers, as well as Marie-Odile Rozon who also served as an observer (Special Advisor for animal transport).

The Committee appreciates the valuable input from the public comment period and all those who provided comments and advice throughout the process.



Summary of Code Requirements

The following is a list of the Requirements within the Dairy Cattle Code of Practice. Refer to the cited Code section for further context about the Requirements.

SECTION 1 Training and Responsibilities

- Personnel must be aware of this Code of Practice and must follow the Requirements of this Code of Practice.
- Personnel must have the competence to carry out the procedures that they are responsible for.
- Managers must supervise personnel and must retrain them if practices begin to fall below standards of care.

SECTION 2 Facilities and Housing

2.1 Design and Maintenance of Facilities

- Housing systems, including flooring and other components of housing, must be maintained in good condition to minimize lameness and injury.
- Electrified crowd gates must not be used.

2.2.1 Calves (Pre-Weaning)

For all calf housing systems:

- Housing must allow calves to easily stand up, lie down, turn completely around, stand fully upright (without touching the top of the enclosure), adopt sternal and lateral resting postures, groom themselves, and have visual contact with other cattle.
- The bedded area for group-housed calves must be large enough to allow all calves to rest comfortably at the same time.
- Where tethering of calves is permitted, the tether must include a collar.

For indoor calf housing:

- Calves must not be tethered as part of normal indoor housing.
- Producers raising calves individually must develop a plan to transition to pair/group housing methods, in consultation with a veterinarian or other qualified advisor.
- Effective April 1, 2031, calves that are healthy, thriving, and compatible must be housed in pairs or groups by 4 weeks of age.¹

Hutches and other outdoor housing:

- Calves housed outdoors, including in hutches, must have physical contact with another calf unless they need to be separated for health reasons or they need to be protected from inclement weather.
- Calves may be tethered only if housed in hutches that provide access to an area outside the hutch.

2.2.2 Heifers

- Housing must allow heifers to easily stand up, lie down, adopt normal resting postures, groom themselves, and have visual and physical contact with other cattle.

2.2.3 Lactating and Dry Cows

- Housing must allow lactating and dry cows to easily stand up, lie down, adopt normal resting postures, groom themselves, and have visual and physical contact with other cattle.
- Effective April 1, 2027, cows must not be tethered continuously throughout their entire production cycle (calving to calving)—they must be provided sufficient regular opportunity for freedom of movement to promote good welfare.

¹ Movement into pairs/groups may need to be delayed for individual calves that are not healthy and thriving. Once moved into pairs/groups, individual calves may need to be singly housed temporarily if they have a health condition that would improve with separation. Movement into pairs/groups may also need to be delayed to ensure there are sufficient number of calves that are compatible as to their age, size, and drinking speed.



Summary of Code Requirements (continued)

- Newly built barns must allow daily, untethered freedom of movement and social interactions year-round.

2.2.4 Breeding Bulls

- Housing must allow breeding bulls to easily stand up, lie down, adopt normal resting postures, groom themselves, turn around, and mount safely.

2.3 Facilities for Special Needs

- Special needs facilities must include a resting surface with bedding that provides comfort, insulation, dryness, and traction.

2.3.1 Calving Areas

- Calving areas, whether for group or individual calving, must provide the cow and calf an area that is clean, safe, and separated from the lactating herd, and that provides enough space for the cow to be assisted.
- Effective April 1, 2029, cattle on all farms must calve in loose housed maternity pens, yards, or pastures that permit them to turn around.
- Newly built barns must allow cows to calve in loose housed maternity pens, yards, or pastures that permit them to turn around.

2.3.3 Areas for Sick, Injured, or Lamé Cattle

- Areas must be available to segregate, care for, and treat cattle that are sick, injured, or lame.

2.4 Ventilation, Temperature, and Relative Humidity

- Facilities, including hutches, must provide cattle with fresh air; prevent the build-up of harmful gases, dust, and moisture; and minimize the risk of heat and cold stress.

2.5 Stall Design

- Stalls and their components must be compatible with the size of the cattle, minimize lameness and injury, and allow cattle to rest comfortably and rise and lie down with ease.
- Tethers or other head restraints must allow cattle to rest in a head back position; stanchions are not permitted.

2.5.1 Electric Trainers in Tie Stalls

- Electric trainers must only be used when needed to train or retrain individual cattle.
- Electric trainers must be safe, secure, adjustable, and positioned to enable normal eating, standing, and lying behaviour.

2.6 Space Allowances

- Stocking density must not exceed 1.2 cows per stall in free stall systems.
- Effective April 1, 2027, stocking density must not normally exceed 1.1 cow per stall.²
- Effective April 1, 2031, stocking density must not normally exceed 1 cow per stall.²
- Resting areas in group pens must provide at least 9.3 m² (100 ft²) per Holstein cow.³

2.7 Feeding Area

- Provide adequate linear feed bunk space to meet the animals' nutritional needs.

² At any time during or after the 2027 and 2031 transition periods, stocking density can go up to 1.2 cows per stall but only temporarily/intermittently.

³ This minimum required space allowance is based on average weights for large breeds (e.g., Holstein) and will be adjusted for medium and small breeds.



Summary of Code Requirements (continued)

2.8 Bedding Management

- Cattle must have a resting surface with bedding that provides comfort, insulation, dryness, and traction.

2.9 Milking Systems

- Milking equipment must be properly maintained and calibrated.

SECTION 3 Feed and Water

3.1 Body Condition Scoring

- Corrective action must be taken for cattle at a body condition score of 2 or lower (refer to *Appendix B – Body Condition Scoring Charts*).

3.2 Nutrition and Feeding Management for Cattle

- Cattle must have daily access to a palatable ration that meets their nutritional needs, promotes satiety, and maintains body condition, health, and vigour.

3.3 Nutrition and Feeding Management for Calves

- Calves must receive a diet that promotes satiety and maintains health, growth, and vigour.
- Newborn calves must be offered a minimum total daily intake of 15% birth weight (6 L for Holsteins) and from 7–28 days of age must be offered a minimum total daily intake of 20% birth weight (8 L for Holsteins) in milk/milk replacer.⁴
- The quantity of milk/milk replacer offered to calves at risk of cold stress must be increased.

3.3.1 Additional Considerations for Weaning

- Calves must be gradually weaned over a period of at least 5 days, and they must be at least 8 weeks old before weaning is completed (44, 48).

3.5 Water

- Watering systems must be clean, and cattle must have access to palatable, clean water in quantities to maintain normal hydration and health, taking into consideration factors such as environmental temperature and diet.
- Neither ice nor snow are suitable sources of water.

SECTION 4 Husbandry Practices

4.1 Handling, Moving, and Restraining Cattle

- Personnel must be knowledgeable in cattle behaviour and must only use low-stress techniques in the routine handling of cattle.
- Electric prods must not be used for routine handling—they must only be used in extreme situations, such as when an animal's safety is at risk.
- Abusive handling is unacceptable.
- When restraint is necessary the safest, least stressful restraint must be used.

4.1.1 Additional Considerations When Handling or Moving Down Cattle

- Apparatus that are designed to lift, move, and support down cattle must be used according to the manufacturer's specifications.

⁴ The amount offered may be reduced for individual calves that are not consistently drinking at this level or for individual calves with health problems.



Summary of Code Requirements (continued)

- Hip lifters must only be used to lift an animal for a short duration to help an animal stand on its own—they must never be used to move down cattle.
- Down cattle must not be moved by hoisting by chain, dragging, or lifting without adequate body support.
- Personnel must not repeatedly encourage a down animal to rise if it has demonstrated it cannot get up or move.
- If an electric prod is used, it must be used in consultation with a veterinarian and only applied on the rear flank and upper rear leg (twice at maximum) when absolutely necessary to determine if the animal can rise or if euthanasia needs to be considered.⁷

4.2 Surgical and Husbandry Procedures

- Surgical procedures must be performed by competent personnel following a method developed in consultation with a veterinarian, including the use of appropriate equipment, pain control, and procedures to minimize the risk of infection and other complications (54, 55).⁷

4.2.1 Animal Identification

- Cattle must not be branded.

4.2.2 Disbudding and Dehorning

- Horn bud removal must be done by 2 months of age (41, 56, 58). Only in exceptional circumstances can individual cattle be dehorned after 2 months of age.
- When removing buds or horns, local anesthesia and systemic analgesia must be provided (4, 55, 58).
- Banding is not an acceptable method of dehorning (59).
- If larger horns must be removed, bleeding must be controlled.

4.2.3 Castration

- If castrating cattle, the procedure must be done as early as possible using local anesthesia and systemic analgesia (4, 55, 60).

4.2.4 Tail Injuries

- Cattle must not be tail docked unless medically necessary for an individual animal, and the procedure must be done using pain control (55).

4.2.5 Extra Teat Removal

- If removing extra teats, they must be removed as early as possible using pain control (55).

4.6 Dry-Off Management

- Cows must not be dried off by restricting water.

SECTION 5 Cattle Health

5.1 Herd Health Management

- Producers must have a veterinarian-client-patient relationship.
- Disease events, treatments, and mortalities (including cause, if known) must be recorded and records must be kept for at least 3 years to track trends in animal health.
- Health records must be reviewed with a veterinarian as part of ongoing herd health and disease prevention planning.

⁷ In consultation with a veterinarian refers to a one-time consultation or periodic consultations as part of a veterinarian-client-patient relationship. Use of this term is not intended to imply that a consultation with a veterinarian is needed each time the procedure or treatment is carried out.



Summary of Code Requirements (continued)

5.1.1 Cattle Cleanliness

- Cattle must be kept clean to minimize disease, maintain udder and hoof health, and promote cattle comfort.

5.3 Caring for Sick, Injured, or Compromised Cattle

- Personnel must be able to detect signs of injury, lameness (including abnormalities in gait or mobility), and disease.
- Sick, injured, lame, or suffering cattle must receive prompt care appropriate to their condition (including pain control where necessary), and they must be monitored at least twice a day.
- Down cattle must have easy access to feed and water and protection from predators, herd mates, and extreme weather (cold, rainfall, direct sunlight), and they must be provided non-slip footing that supports recovery.
- Lactating cows that are severely lame or down that require milking (to prevent mammary engorgement) must be milked where they are located.

5.4 Calving Management

- Steps must be taken to ensure cattle calve in a designated calving area.
- Cattle close to calving must be monitored daily, at intervals suitable to individual cattle needs, including for risk factors of dystocia.

5.5 Cattle Health

- If mortality in female calves from 2 days of age exceeds 10%, corrective actions must be implemented to improve calving management and calf health in consultation with a veterinarian or other qualified advisor (77).

5.5.1 Colostrum

- Male and female calves must receive at least 4 liters of good quality colostrum within 12 hours of birth, with the first meal occurring as soon as possible, and no later than 6 hours after birth (41).

5.6 Preventing and Treating Mastitis

- Systemic analgesia must be included in the treatment of cows with severe acute clinical mastitis (4).

5.7 Promoting Optimal Foot and Leg Health

- Personnel must regularly observe cattle for signs of lameness or leg injuries, to diagnose and treat them quickly.
- To minimize lameness and leg injuries, producers must set thresholds for the occurrence of lameness and leg injuries and take corrective actions when the thresholds are exceeded.

5.7.1 Hoof Trimming

- Feet and claws must be inspected and hooves trimmed as required to promote a normal gait and minimize lameness.
- Infectious hoof lesions must be treated to control the infection.
- Therapeutic hoof trimming must include strategies to relieve pain and pressure on the injured area and promote healing.
- Pain control must be included in the treatment of cattle that receive an invasive hoof trim.



Summary of Code Requirements (continued)

SECTION 6 Preparations for Transport

6.1.1 Fitness for Transport (General and Cull Cows)

- Every animal's fitness for transport must be assessed before they are loaded taking into consideration their condition and risk factors that may impact the animal's capacity to withstand loading, transport, and unloading (3).
- Unfit animals must not be transported except for veterinary care on the advice of a veterinarian and with special provisions (refer to the regulatory guidance in *Appendix F – Transport Decision Tree*) (3).¹⁰
- Compromised animals must only be transported with special provisions and directly to the nearest suitable place where they can receive care or be promptly slaughtered (not through an auction or assembly yard; refer to the regulatory guidance in *Appendix F – Transport Decision Tree*) (3).¹⁰

6.1.2 Additional Considerations for Calves

- Calves must only be shipped if they are free from signs of fever or disease and have a healed, uninfected navel (refer to other regulatory guidance in *Appendix F – Transport Decision Tree*).¹¹
- Calves that are 8 days of age or less must only be transported with special provisions and directly to their final destination (not through an auction or assembly yard) (3).¹²

6.1.3 Preparing Cattle for Transport

- When shipping a lactating cow, steps must be taken to prevent the risk of udder engorgement.

6.1.4 Arranging Transport

- Personnel involved in loading, unloading, or transporting cattle must have the necessary knowledge and skills to conduct these activities in compliance with the Health of Animals Regulations (3).

6.2 Loading and Unloading

- Ramps, gangways, chutes, and steps must be designed, constructed, maintained, and used in a way that prevents the animal from tripping, slipping, or falling (3).
- The slope of ramps used to load or unload animals onto/from the conveyance must not exceed 25 degrees (3).
- Cattle that are incompatible must be segregated (3).

SECTION 7 Euthanasia

7.1 Decision Making and Criteria for Euthanasia

- Cattle must be promptly euthanized if they have a condition that compromises their welfare AND at least 1 of the following applies:
 - they do not have a reasonable prospect of improvement,
 - they are not responding to treatment(s) within an appropriate timeframe for the condition, or
 - treatment is not a humane option.
- Personnel who perform euthanasia must be trained and competent in all aspects of the acceptable method(s) used on the farm.

¹⁰ The *Health of Animals Regulations* require that compromised or unfit animals are individually loaded and unloaded without having to negotiate ramps inside the conveyance, are isolated during transport (compromised animals may be segregated with one familiar animal), and other measures are taken as necessary to prevent suffering or injury (e.g., mitigate pain, protect from cold, prevent dehydration).

¹¹ From the *Health of Animals Regulations*: calves with signs of fever or that have an unhealed or infected navel are unfit and must only be transported for veterinary care on the advice of a veterinarian with special provisions.

¹² The *Health of Animals Regulations* require that calves that are 8 days of age or less are individually loaded and unloaded without having to negotiate ramps inside the conveyance; have sufficient space to allow the animal to lie down without lying on top of another animal; are segregated from older animals (other than their dam); and that other measures are taken as necessary to prevent suffering, injury, or death.



Summary of Code Requirements (continued)

7.2 Methods

- An acceptable method for euthanizing cattle must be used (acceptable methods are listed in Table 7.1).
- The method of euthanasia must be quick, cause minimal stress and pain, and result in rapid loss of consciousness followed by death without the animal regaining consciousness.
- Manual blunt force trauma is not an acceptable means of euthanasia, including for calves at any age (5, 100, 101).
- Every farm must have the ability to euthanize animals or have timely access to euthanasia services.
- Equipment necessary for euthanasia (including a secondary step, if applicable) must be used, stored, and maintained according to the manufacturer's instructions to ensure proper function.
- Prior to being euthanized, cattle must only be handled or moved if necessary.
- When restraint is necessary, euthanasia must be performed without delay following restraint.

7.3 Confirming Loss of Consciousness and Death

- Cattle euthanized using gunshot or euthanasia drugs must be assessed to confirm that they are unconscious immediately after application. If an animal is not immediately unconscious, then a repeat application must be immediately delivered.
- Cattle euthanized using a penetrating captive bolt must be assessed to confirm that they are unconscious immediately after application and before the secondary step to cause death is applied. If an animal is not immediately unconscious, then a repeat application must be immediately delivered.
- Before moving or leaving the animal, death must be confirmed.

